# antibodies -online.com







## anti-CD93 antibody (FITC)

**Images** 



#### Overview

Quantity:	100 tests
Target:	CD93
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD93 antibody is conjugated to FITC
Application:	Flow Cytometry (FACS)


## **Product Details**


Immunogen:	KG1 cell line
Clone:	VIMD2
Isotype:	IgG1
Specificity:	The mouse monoclonal antibody VIMD2 recognizes an extracellular epitope on CD93, an approximately 110-120 kDa glycoprotein expressed mainly on myeloid cells and endothelial cells.
Purification:	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

## Target Details

## **Target Details**

3.7		
Alternative Name:	CD93 (CD93 Products)	
Background:	CD93 Molecule,CD93 (also known as C1qR1) is a type I transmembrane glycoprotein	
	containing extracellular N-terminal C-type lectin domain and five EGF-like domains, and an	
	intracellular tail interacting with moesin, a protein known to play a role in linking	
	transmembrane proteins to the cytoskeleton and in the remodelling of the cytoskeleton. CD93	
	was reported to serve as a receptor for complement component C1q, but this function has not	
	been fully elucidated yet. CD93 is involved in intercellular adhesion and in the clearance of	
	apoptotic cells.,C1QR1, C1qRP, ECSM3, MXRA4, C1qR(P), dJ737E23.1	
Gene ID:	22918	
UniProt:	Q9NPY3	
Application Details		
Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µL reagent /	
	100 $\mu L$ of whole blood or $10^6$ cells in a suspension. The content of a vial (0.4 ml) is sufficient for	
	100 tests.	
Restrictions:	For Research Use only	
Handling		
Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Storage:	4 °C	
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.	





### **Flow Cytometry**

**Image 1.** Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD93 (VIMD2) FITC antibody (4  $\mu$ L reagent / 100  $\mu$ L of peripheral whole blood).

#### **Flow Cytometry**

**Image 2.** Separation of human monocytes (red-filled) from lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD93 (VIMD2) FITC antibody ( $4 \mu L$  reagent /  $100 \mu L$  of peripheral whole blood).