.-online.com antibodies

Datasheet for ABIN7013978 anti-TNFRSF13C antibody

2 Images



Overview

Quantity:	0.1 mg
Target:	TNFRSF13C
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TNFRSF13C antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	human CD268-transfected murine L1.2 cells
Clone:	11C1
lsotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody 11C1 recognizes an extracellular epitope of CD268 / BAFF R (B cell-activating factor receptor), a 19 kDa type III transmembrane protein expressed on resting B cells and CD4-positive T cells, but down regulated after activation.
Purification:	Purified by protein-A affinity chromatography.
Target Details	

Target:	TNFRSF13C
Alternative Name:	CD268 (TNFRSF13C Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7013978 | 03/05/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Background:	TNF receptor superfamily member 13C,CD268 / BAFF R is a TNFR family receptor that binds
	the B-cell-activating factor (CD257 / BAFF). Splice variants of CD268 have been observed both
	in man and mouse. A naturally occurring mutation of CD268 in A/WySnJ mice is associated
	with low number of mature B cells, but with normal B cell precursors. The role of BAFF in B-cell
	survival and activation make CD268 a potential diagnostic reagent. It may be involved in
	survival of B-cell malignancies. Experimental administration of a CD268-Fc fusion protein
	suppresses antibody responses. In T cells the CD268 costimulates their activation and
	proliferation. Defects in CD268 cause the common variable immunodeficiency 4
	(CVID4).,TNFRSF13C, BAFFR, CVID4, BAFF-R, BROMIX, prolixin


Gene ID:	115650
UniProt:	Q96RJ3
Pathways:	NF-kappaB Signaling

Application Details

Application Notes:	Flow cytometry: Recommended dilution: 1-4 µg/mL.
Restrictions:	For Research Use only

Handling

Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.



Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD268 (11C1) purified antibody (concentration in sample $0,6 \mu$ g/mL, GAM APC).

Flow Cytometry

Image 2. Separation of human CD268 positive lymphocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD268 (11C1) purified antibody (concentration in sample $0,6 \mu$ g/mL, GAM APC).

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7013978 | 03/05/2024 | Copyright antibodies-online. All rights reserved.