antibodies -online.com





anti-TNFRSF13C antibody (PE)

2 Images



Go to Product page

Overview

Quantity:	100 tests
Target:	TNFRSF13C
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TNFRSF13C antibody is conjugated to PE
Application:	Flow Cytometry (FACS)

Product Details


Immunogen:	human CD268-transfected murine L1.2 cells
Clone:	11C1
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody 11C1 recognizes an extracellular epitope of CD268 / BAFF R (B cell-activating factor receptor), a 19 kDa type III transmembrane protein expressed on resting B cells and CD4-positive T cells, but down regulated after activation.
Purification:	Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.


Target Details

Target:	TNFRSF13C
Alternative Name:	CD268 (TNFRSF13C Products)

Target Details

-	
Background:	TNF receptor superfamily member 13C,CD268 / BAFF R is a TNFR family receptor that binds
	the B-cell-activating factor (CD257 / BAFF). Splice variants of CD268 have been observed both
	in man and mouse. A naturally occurring mutation of CD268 in A/WySnJ mice is associated
	with low number of mature B cells, but with normal B cell precursors. The role of BAFF in B-cell
	survival and activation make CD268 a potential diagnostic reagent. It may be involved in
	survival of B-cell malignancies. Experimental administration of a CD268-Fc fusion protein
	suppresses antibody responses. In T cells the CD268 costimulates their activation and
	proliferation. Defects in CD268 cause the common variable immunodeficiency 4
	(CVID4).,TNFRSF13C, BAFFR, CVID4, BAFF-R, BROMIX, prolixin
Gene ID:	115650
UniProt:	Q96RJ3
Pathways:	NF-kappaB Signaling
Application Details	
Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µL reagent
	/ 100 μL of whole blood or 10^6 cells in a suspension. The content of a vial (1 ml) is sufficient for
	100 tests.
Restrictions:	For Research Use only
Handling	
Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.





Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD268 (11C1) PE antibody (10 μ L reagent / 100 μ L of peripheral whole blood).

Flow Cytometry

Image 2. Separation of human CD268 positive lymphocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD268 (11C1) PE antibody (10 μ L reagent / 100 μ L of peripheral whole blood).