

Datasheet for ABIN7013986

anti-CD21 antibody (PerCP)





Go to Product page

_						
	1//	Д	rv	16	٦/	٨
	W	\vdash	ΙV	Ιt	٦,	/V


Quantity:	100 tests	
Target:	CD21 (CR2)	
Reactivity:	Human, Pig, Cow, Dog	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This CD21 antibody is conjugated to PerCP	
Application:	Flow Cytometry (FACS)	

Product Details

Purpose:	Anti-Hu CD21 PerCP
Immunogen:	IM9 human B-lymphoblastoid cell line
Clone:	LT21
Isotype:	IgG1 kappa
Specificity:	The antibody LT21 reacts with an extracellular epitope of CD21 (CR2), a 145 kDa transmembrane glycoprotein (complement C3d receptor - C3dR) expressed on B lymphocytes, follicular dendritic cells, some epithelial cells and a subsets of T lymphocytes. It is not expressed on immature B cells.
Purification:	Purified antibody is conjugated with activated Peridinin-Chlorophyll Protein (PerCP) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Target Details

Target:	CD21 (CR2)		
Alternative Name:	CD21 (CR2 Products)		
Background:	Complement C3d receptor 2,CD21 (complement receptor 2, CR2) binds C3 complement fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of immunocomplexes, survival of memory B cells and in development and maintenance of the humoral response to T-dependent antigens. CD21 also serves as a key receptor for Epstein-Barrius binding and is involved in targeting prions to folicular dendritic cells and expediting neuroinvasion following peripheral exposure to prions. A soluble form of the CD21 (sCD21) is		
	shed from the lymphocyte surface and retains its ability to bind respective ligands.,CR2, C3DR2, CVID7, SLEB9		
Gene ID:	1380		
UniProt:	P20023		
Pathways:	Complement System		
Application Details			
Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 10 μ L reagent / 100 μ L of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.		
Restrictions:	For Research Use only		
Handling			
Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.		
Storage:	4 °C		
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.		



Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD21 (LT21) PerCP antibody (10 μ L reagent / 100 μ L of peripheral whole blood).

Flow Cytometry

Image 2. Separation of human CD21 lymphocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD21 (LT21) APC antibody (10 µL reagent / 100 µL of peripheral whole blood).