antibodies -online.com







anti-F11R antibody (AA 51-150)





\sim							
	1//	\Box	r۱	/ [\bigcirc	1	٨,

Quantity:	100 μL
Torquet	F11D
Target:	F11R
Binding Specificity:	AA 51-150
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This F11R antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Junctional Adhesion Molecule 1
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Pig
Purification:	Purified by Protein A.

Target Details

Target:	F11R
Alternative Name:	JAM1 (F11R Products)
Background:	Synonyms: JAM, KAT, JAM1, JAMA, JCAM, CD321, PAM-1, Junctional adhesion molecule A,

JAM-A, Junctional adhesion molecule 1, JAM-1, Platelet F11 receptor, Platelet adhesion
molecule 1, F11R, UNQ264/PRO301

Background: Seems to play a role in epithelial tight junction formation. Appears early in primordial forms of cell junctions and recruits PARD3. The association of the PARD6-PARD3 complex may prevent the interaction of PARD3 with JAM1, thereby preventing tight junction assembly (By similarity). Plays a role in regulating monocyte transmigration involved in integrity of epithelial barrier. Involved in platelet activation. In case of orthoreovirus infection, serves as receptor for the virus.

Gene ID:	50848
UniProt:	Q9Y624

Cell-Cell Junction Organization

Application Details

Restrictions:	For Research Use only
	ELISA 1:500-1000
Application Notes:	WB 1:300-5000

Handling

Pathways:

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



Immunohistochemistry

Image 1. Formalin-fixed and paraffin embedded mouse small intestine tissue labeled with Anti JAM1 Polyclonal Antibody, Unconjugated (ABIN702130) at 1:200 followed by conjugation to the secondary antibody and DAB staining