## antibodies -online.com





Datasheet for ABIN702465 anti-DIO2 antibody (AA 101-200) (Cy5.5)

Go to Product page

| ( ) | ۱ ۱ | $\cap$ | r  | /1 | $\cap$ | ۱ ۸ | 1 |
|-----|-----|--------|----|----|--------|-----|---|
| 0   | 'V  | ㄷ      | I١ | νı | ㄷ      | ٧   | ۷ |

| Quantity:            | 100 μL   |  |
|----------------------|--|--|
| Target:              | DIO2   |  |
| Binding Specificity: | AA 101-200   |  |
| Reactivity:          | Mouse, Rat   |  |
| Host:                | Rabbit   |  |
| Clonality:           | Polyclonal   |  |
| Conjugate:           | This DIO2 antibody is conjugated to Cy5.5  |  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |

## **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human DIO2 |  |
|-----------------------|--|--|
| Isotype:              | IgG  |  |
| Cross-Reactivity:     | Mouse, Rat   |  |
| Predicted Reactivity: | Human,Dog,Cow,Sheep,Pig,Horse,Rabbit                     |  |
| Purification:         | Purified by Protein A.                                   |  |

## **Target Details**

| Target:           | DIO2                 |
|-------------------|----------------------|
| Alternative Name: | DIO2 (DIO2 Products) |

## **Target Details**

| Background:         | Synonyms: 5DII, D2, Deiodinase, iodothyronine, type II, DIOII, ITDI2, SelY, Thyroxine deiodinase, type II, TXDI2, Type 2 DI, Type II 5' deiodinase, Type II iodothyronine deiodinase.  Background: DIO2 belongs to the iodothyronine deiodinase family and is responsible for the deiodination of T4 (3,5,3',5'-tetraiodothyronine) into T3 (3,5,3'-triiodothyronine). It is essential for providing the brain with appropriate levels of T3 during the critical period of development. DIO2 is expressed in heart, skeletal muscle, placenta, fetal brain and several regions of the adult |
|---------------------|---|
|                     | brain. There are two named isoforms.  |
| Gene ID:            | 1734  |
| Pathways:           | Hormone Transport, Hormone Activity   |
| Application Details |   |
| Application Notes:  | IF(IHC-P) 1:50-200  |
|                     | IF(IHC-F) 1:50-200  |
|                     | IF(ICC) 1:50-200  |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Concentration:      | 1 μg/μL   |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.  |
| Preservative:       | ProClin   |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.  |
| Storage:            | -20 °C  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.   |
| Expiry Date:        | 12 months   |