

Datasheet for ABIN702942

anti-p53 antibody (pSer37) (Biotin)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	p53 (TP53)
Binding Specificity:	pSer37
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This p53 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human P53 around the phosphorylation site of Ser37
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

Target:	p53 (TP53)
Alternative Name:	P53 (TP53 Products)

Target Details

Background:	Synonyms: p53Ser37, P53 phospho S37, P53 phospho Ser37, p-P53 Ser37, p-P53 S37, Widespread p53, Wtp53, Antigen NY-CO-13, Cellular tumor antigen p53, Cys 51 Stop, HGNC11998, LFS1, p53, p53 Cellular Tumor Antigen, p53 Tumor Suppressor, Phosphoprotein p53, TP53, Transformation related protein 53, TRP53, Tumor protein p53, Tumour Protein p53, P53_HUMAN, TP53. Background: This gene encodes a tumor suppressor protein containing transcriptional activation, DNA binding, and oligomerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni syndrome. Alternative splicing of this gene and the use of alternate promoters result in multiple transcript variants and isoforms. Additional isoforms have also been shown to result from the use of alternate translation initiation codons (PMIDs: 12032546, 20937277). [provided by RefSeq, Feb 2013].
Gene ID:	7157
Pathways:	p53 Signaling , MAPK Signaling , PI3K-Akt Signaling , Apoptosis , AMPK Signaling , Chromatin Binding , ER-Nucleus Signaling , Positive Regulation of Endopeptidase Activity , Hepatitis C , Protein targeting to Nucleus , Autophagy , Warburg Effect

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months