

Datasheet for ABIN703000
anti-p53 antibody (pThr18)



[Go to Product page](#)

1 Publication

Overview

Quantity:	100 µL
Target:	p53 (TP53)
Binding Specificity:	pThr18
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This p53 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human P53 around the phosphorylation site of Thr18
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	p53 (TP53)
Alternative Name:	P53 (TP53 Products)
Background:	Synonyms: p53 Thr18, P53 phospho T18, P53 phospho Thr18, p-P53 Thr18, p-P53 T18, Widespread p53, Wtp53, Antigen NY-CO-13, Cellular tumor antigen p53, Cys 51 Stop, HGNC11998, LFS1, p53, p53 Cellular Tumor Antigen, p53 Tumor Suppressor, Phosphoprotein p53, TP53, Transformation related protein 53, TRP53, Tumor protein p53, Tumour Protein p53, P53_HUMAN, TP53. Background: This gene encodes a tumor suppressor protein containing transcriptional activation, DNA binding, and oligomerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni syndrome. Alternative splicing of this gene and the use of alternate promoters result in multiple transcript variants and isoforms. Additional isoforms have also been shown to result from the use of alternate translation initiation codons (PMIDs: 12032546, 20937277). [provided by RefSeq, Feb 2013].
Gene ID:	7157
Pathways:	p53 Signaling , MAPK Signaling , PI3K-Akt Signaling , Apoptosis , AMPK Signaling , Chromatin Binding , ER-Nucleus Signaling , Positive Regulation of Endopeptidase Activity , Hepatitis C , Protein targeting to Nucleus , Autophagy , Warburg Effect

Application Details

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 FCM 1:20-100 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

Publications

Product cited in:	He, Zeng, Yu, Zhang, Zeng, Gong, Duan, Liu, Yang: "Resveratrol Improved the Progression of Chronic Prostatitis via the Downregulation of c-kit/SCF by Activating Sirt1." in: Journal of agricultural and food chemistry , Vol. 65, Issue 28, pp. 5668-5673, (2017) (PubMed).
-------------------	---