

Datasheet for ABIN703017 anti-p53 antibody (pThr81) (Biotin)



Overview

Overview	
Quantity:	100 μL
Target:	p53 (TP53)
Binding Specificity:	pThr81
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This p53 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)),
	Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human P53 around the
	phosphorylation site of Thr81
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Chicken
Purification:	Purified by Protein A.
Target Details	
Target:	p53 (TP53)

Target Details

Alternative Name:	P53 (TP53 Products)
Background:	Synonyms: p53 Thr81, P53 phospho T81, P53 phospho Thr81, p-P53 Thr81, p-P53 T81,
	Widespread p53, Wtp53, Antigen NY-CO-13, Cellular tumor antigen p53, Cys 51 Stop,
	HGNC11998, LFS1, p53, p53 Cellular Tumor Antigen, p53 Tumor Suppressor, Phosphoprotein
	p53, TP53, Transformation related protein 53, TRP53, Tumor protein p53, Tumour Protein p53, P53_HUMAN, TP53.
	Background: This gene encodes a tumor suppressor protein containing transcriptional
	activation, DNA binding, and oligomerization domains. The encoded protein responds to diverse
	cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest,
	apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are
	associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni
	syndrome. Alternative splicing of this gene and the use of alternate promoters result in multiple
	transcript variants and isoforms. Additional isoforms have also been shown to result from the
	use of alternate translation initiation codons (PMIDs: 12032546, 20937277). [provided by
	RefSeq, Feb 2013].
Gene ID:	7157
Pathways:	p53 Signaling, MAPK Signaling, PI3K-Akt Signaling, Apoptosis, AMPK Signaling, Chromatin
	Binding, ER-Nucleus Signaling, Positive Regulation of Endopeptidase Activity, Hepatitis C,
	Protein targeting to Nucleus, Autophagy, Warburg Effect
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months