

Datasheet for ABIN703075

anti-TCP1 alpha/CCTA antibody (pSer455)



Overview

| Overview | |
|----------------------|--|
| Quantity: | 100 μL |
| Target: | TCP1 alpha/CCTA (TCP1) |
| Binding Specificity: | pSer455 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This TCP1 alpha/CCTA antibody is un-conjugated |
| Application: | ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |
| Product Details | |
| Immunogen: | KLH conjugated synthetic phosphopeptide derived from human p63 around the phosphorylation site of Ser455 |
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Purification: | Purified by Protein A. |
| Target Details | |
| Target: | TCP1 alpha/CCTA (TCP1) |
| Alternative Name: | P63 (TCP1 Products) |
| | |

Target Details

Background:

Synonyms: p63 phospho S455, p63 phospho Ser455, p-p63 Ser455, AIS, Amplied in squamous cell carcinoma, Bp51A, Bp51B, p63 Alpha, Chronic ulcerative stomatitis protein, CUSP, DN p63 alpha 1, DNp63, EEC3, Keratinocyte transcription factor, Keratinocyte transcription factor KET, KET, LMS, NBP, OFC8, p40, p51, P51/P63, p53 like transcription factor, p53 related protein, p53related protein p63, p53CP, p63, P73, p73H, p73L, RHS, SHFM4, TP53CP, TP53L, TP63, TP73, TP73L, Transformation related protein 63, Trp53rp1, Trp63, Tumor protein 63, tumor protein 63 kDa with strong homology to p53, Tumor protein p53-competing protein, Tumor protein p53like, tumor protein p63, Tumor protein p73, tumor protein p73-like, P63_HUMAN. Background: This gene encodes a member of the p53 family of transcription factors. An animal model, p63 -/- mice, has been useful in defining the role this protein plays in the development and maintenance of stratified epithelial tissues. p63 -/- mice have several developmental defects which include the lack of limbs and other tissues, such as teeth and mammary glands, which develop as a result of interactions between mesenchyme and epithelium. Mutations in this gene are associated with ectodermal dysplasia, and cleft lip/palate syndrome 3 (EEC3), split-hand/foot malformation 4 (SHFM4), ankyloblepharon-ectodermal defects-cleft lip/palate, ADULT syndrome (acro-dermato-ungual-lacrimal-tooth), limb-mammary syndrome, Rap-Hodgkin syndrome (RHS), and orofacial cleft 8. Both alternative splicing and the use of alternative promoters results in multiple transcript variants encoding different proteins. Many transcripts encoding different proteins have been reported but the biological validity and the full-length nature of these variants have not been determined. [provided by RefSeq, Jul 2008].

Gene ID:

8626

Application Details

Application Notes:

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:

Liquid

Concentration:

 $1 \mu g/\mu L$

Handling

| Buffer: | 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. |
|--------------------|--|
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
| Expiry Date: | 12 months |