# antibodies -online.com







# anti-DDX19B antibody (AA 51-150) (Biotin)



| ( ) | ve     | K\ / |    | A . |
|-----|--------|------|----|-----|
|     | $\cup$ | 1 V/ | Щ. | V۷  |
|     |        |      |    |     |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | DDX19B   |
| Binding Specificity: | AA 51-150  |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This DDX19B antibody is conjugated to Biotin   |
| Application:         | ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

## **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human DDX19B |
|-----------------------|--|
| Isotype:              | IgG  |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse                    |
| Purification:         | Purified by Protein A.                                     |

# **Target Details**

| Target:   | DDX19B                   |
|---|--------------------------|
| Alternative Name:   | DDX19B (DDX19B Products) |
| Background: Synonyms: DDX19B, Dbp5, DDX19, DEAD box protein 19B, DEAD box RNA helicase DE |                          |

helicase DEAD5, DD19B\_HUMAN, ATP-dependent RNA helicase DDX19B, DEAD box RNA helicase DEAD5, DBP5, TDBP.

Background: DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein, which exhibits RNA-dependent ATPase and ATP-dependent RNA-unwinding activities. This protein is recruited to the cytoplasmic fibrils of the nuclear pore complex, where it participates in the export of mRNA from the nucleus. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008].

Gene ID:

11269

#### **Application Details**

Application Notes: IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

### Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C for 12 months.  |
| Expiry Date:       | 12 months  |