

Datasheet for ABIN7043701

anti-GLUT4 antibody (Intracellular) (Atto 594)



[Go to Product page](#)

1 Image

Overview

Quantity:	50 µL
Target:	GLUT4 (SLC2A4)
Binding Specificity:	AA 495-507, Intracellular
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GLUT4 antibody is conjugated to Atto 594
Application:	Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Purpose:	A Rabbit Polyclonal Antibody to Glucose Transporter 4 Conjugated to the Fluorescent Dye ATTO-594
Immunogen:	Immunogen: Synthetic peptide Immunogen Sequence: (C)KPSTELEYLGPDE, corresponding to amino acid residues 495-507 of human Glucose Transporter 4
Isotype:	IgG
Specificity:	Intracellular, C-terminus
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Rat,mouse - identical
Characteristics:	Anti-GLUT4 Antibody (ABIN7043700, ABIN7044487 and ABIN7044488) is a highly specific

antibody directed against an epitope of the human Glucose Transporter 4 protein. The antibody can be used in western blot and immunohistochemistry applications. It has been designed to recognize GLUT4 from human, rat, and mouse samples. \nAnti-GLUT4-ATTO Fluor-594 Antibody (ABIN7043701) is directly labeled with an fluorescent dye. ATTO dyes are characterized by strong absorption (high extinction coefficient), high fluorescence quantum yield, and high photo-stability. The fluorescent label belongs to the class of Rhodamine dyes and can be used with fluorescent equipment typically optimized to detect Texas Red and Alexa-594. Anti-GLUT4-ATTO Fluor-594 Antibody is especially suited for experiments requiring simultaneous labeling of different markers.

Purification: Affinity purified on immobilized antigen.

Target Details

Target: GLUT4 (SLC2A4)

Alternative Name: SLC2A4 ([SLC2A4 Products](#))

Background: Glucose transporter 4, Facilitated glucose transporter member 4, Insulin-responsive glucose transporter type 4, SLC2A4,The GLUT protein family is a member of the major facilitator superfamily of membrane transporters and is encoded by the SLC2 genes. GLUT proteins contain 12 membrane-spanning domains, an N-linked glycosylation site, and intracellular NH2 and COOH termini. In addition, there are several conserved residues and motifs designated "sugar transporter signatures".To date, 14 GLUT isoforms have been identified and divided into 3 different classes based on sequence similarity and structural and functional characteristics. GLUT4 together with GLUT 1-3 and 14 belong to Class I.GLUT4 is an insulin-regulated glucose transporter responsible for insulin-regulated glucose uptake into fat and muscle cells. GLUT4 translocates from intracellular stores to the cell surface, from an inactive site to an active one in order to enhance glucose uptake1.GLUT4 is expressed mainly in adipose tissue and skeletal muscle. It is also detected in brain, kidney and intestine.There is no evidence for mutation in GLUT4 that is linked to metabolic disease in humans. However, GLUT4 polymorphisms are rare and equally common in normoglycaemic individuals and type 2 diabetics. General decrease in GLUT4 levels show signs of diabetes, while a disruption in GLUT4 expression in muscle or adipose tissue might contribute to the development of whole-body insulin resistance in humans2.

Alternative names: GLUT4, Glucose transporter 4, Facilitated glucose transporter member 4, Insulin-responsive glucose transporter type 4, SLC2A4

Target Details


Gene ID:	6517
NCBI Accession:	NM_001042
UniProt:	P14672
Pathways:	AMPK Signaling , Carbohydrate Homeostasis , Proton Transport , Brown Fat Cell Differentiation , Warburg Effect

Application Details

Application Notes:	Antigen preadsorption control: 1 µg peptide per 1 µg antibody Application Dilutions Immunohistochemistry paraffin embedded sections ihc: 1:40 Application Dilutions Western blot wb: N/A
Comment:	Negative Control: (ABIN7237006) Blocking Peptide: (ABIN7237006)
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	50 µL double distilled water (DDW).
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 1 % BSA with 0.05 % sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Storage before reconstitution: The antibody ships as a lyophilized powder at room temperature. Upon arrival, it should be stored at -20°C. Storage after reconstitution: The reconstituted solution can be stored at 4°C, protected from the light, for up to 1 week. For longer periods, small aliquots should be stored at -20°C. Avoid multiple freezing and thawing. Centrifuge all antibody preparations before use (10000 x g 5 min).



Immunohistochemistry

Image 1. Expression of Glucose Transporter 4 in rat olfactory bulb - Immunohistochemical staining of rat olfactory bulb free floating frozen sections using Anti-GLUT4-ATTO Fluor-594 Antibody (ABIN7043701), (1:40). GLUT4 immunoreactivity (red) is strongly detected in some glomeruli (vertical arrows). Lower immunoreactivity in other glomeruli (horizontal arrows) is also observed. Nuclei are stained using DAPI as the counterstain.