antibodies - online.com







anti-TRPV2 antibody (1st Extracellular Loop)



Images



| \sim | | | |
|--------|-----------------------|------|---------------|
| | $ \backslash / \cap$ | r\/I | $\triangle V$ |


| Quantity: | 25 μL | |
|----------------------|---|--|
| Target: | TRPV2 | |
| Binding Specificity: | 1st Extracellular Loop, AA 413-428 | |
| Reactivity: | Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP), Immunocytochemistry (ICC), Flow Cytometry (FACS), Live Cell Imaging (LCI) | |

Product Details

| Immunogen: | Immunogen: Synthetic peptide Immunogen Sequence: (C)HQPSLDQPAIPSSKAT, corresponding to amino acid residues 413- 428 of rat TRPV2 |
|------------------|--|
| Isotype: | IgG |
| Characteristics: | Anti-TRPV2 (VRL1) (extracellular) Antibody (ABIN7043845, ABIN7044002 and ABIN7044003)) is a highly specific antibody directed against an epitope of the rat protein. The antibody can be used in western blot, immunoprecipitation, immunohistochemistry, live cell imaging, and indirect flow cytometry applications. It has been designed to recognize TRPV2 from mouse and rat samples. |
| Purification: | Affinity purified on immobilized antigen. |


Target Details

| Target: | TRPV2 |
|--|--|
| Alternative Name: | TRPV2 (VRL1) (TRPV2 Products) |
| Background: | Alternative names: TRPV2 (VRL1), Transient receptor potential cation channel subfamily V member 2, Vanilloid receptor-like protein 1, VRL-1, Osm-9-like TRP channel 2, OTRPC2, Stretch-activated channel 2B, SAC2B |
| Gene ID: | 29465 |
| NCBI Accession: | NM_016113 |
| UniProt: | Q9WUD2 |
| Pathways: | Regulation of Cell Size |
| Application Details | |
| Application Notes: | Optimal working dilution should be determined by the investigator. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Lyophilized |
| | |
| Reconstitution: | $25\mu\text{L}$, $50\mu\text{L}$ or 0.2mL double distilled water (DDW), depending on the sample size. |
| Reconstitution: Concentration: | $25\mu\text{L}, 50\mu\text{L}$ or 0.2 mL double distilled water (DDW), depending on the sample size. |
| | |
| Concentration: | 0.6 mg/mL Reconstituted antibody contains phosphate buffered saline (PBS), pH 7.4, 1 % BSA, 0.05 % |
| Concentration: Buffer: | 0.6 mg/mL Reconstituted antibody contains phosphate buffered saline (PBS), pH 7.4, 1 % BSA, 0.05 % Sodium azide. |
| Concentration: Buffer: Preservative: | 0.6 mg/mL Reconstituted antibody contains phosphate buffered saline (PBS), pH 7.4, 1 % BSA, 0.05 % Sodium azide. Sodium azide This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which |




Western Blotting

Image 1. Western blot analysis of ND7/23 cell line membrane (lanes 1 and 3), RBL lysates (lanes 2 and 4) and rat brain membrane (lanes 5 and 6): - 1,2,5. Anti-TRPV2 (VRL1) (extracellular) Antibody (ABIN7043845, ABIN7044002 and ABIN7044003), (1:200).3,4,6. Anti-TRPV2 (VRL1) (extracellular) Antibody, preincubated with TRPV2/VRL1 (extracellular) Blocking Peptide (#BLP-CC039).



Immunohistochemistry

Image 2. Expression of TRPV2 in mouse DRG - Immunohistochemical staining of TRPV2 in mouse dorsal root ganglion (DRG) using Anti-TRPV2 (VRL1) (extracellular) Antibody (ABIN7043845, ABIN7044002 and ABIN7044003). A. TRPV2 (green) appears in patches along the perimeter of the DRG (arrows). B. Neurons containing neurofilament 200 (red) are scattered in the DRG, also in patches (arrows). C. A merge of the two panels shows that the spatial distribution of neurofilament 200 and TRPV2 expression overlaps. However, DRGs showing robust expression of neurofilament 200 do not contain TRPV2.



Immunoprecipitation

Image 3. Immunoprecipitation of rat basophilic leukemia (RBL) cell lysate: - 1. RBL lysate.2. Lysate with Anti-TRPV2 immunoprecipitated (VRL1) (extracellular) Antibody (ABIN7043845, ABIN7044002 and ABIN7044003), (6 µg).3. Lysate immunoprecipitated with pre-immune rabbit serum. The upper arrow indicates TRPV2 while the lower arrow indicates the IgG heavy chain. Western blot analysis was performed with Anti-TRPV2 (VRL1) (extracellular) Antibody.

Please check the product details page for more images. Overall 5 images are available for ABIN7043845.