

## Datasheet for ABIN704420 anti-PLAA antibody (AA 101-200) (HRP)



| Overview              |   |
|-----------------------|---|
| Quantity:             | 100 μL  |
| Target:               | PLAA  |
| Binding Specificity:  | AA 101-200  |
| Reactivity:           | Human   |
| Host:                 | Rabbit  |
| Clonality:            | Polyclonal  |
| Conjugate:            | This PLAA antibody is conjugated to HRP   |
| Application:          | Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |
|                       |   |
| Product Details       |   |
| Immunogen:            | KLH conjugated synthetic peptide derived from human Phospholipase A2 activator protein  |
| Isotype:              | IgG   |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit  |
| Purification:         | Purified by Protein A.  |
| Target Details        |   |
| Target:               | PLAA  |
| Alternative Name:     | PLAA/Phospholipase A2 activator protein (PLAA Products)   |
| Background:           | Synonyms: DOA1, PLAP, PLA2P, Phospholipase A-2-activating protein, PLAA   |

| Background: Phospholipase A2 activator protein (PLAP), activates PLA2 (phospholipase A2           |  |
|---|--|
| enzyme) and is an important mediator of eicosanoid generation. Also known as a pro-               |  |
| inflammatory agent, PLAP has been found in inflamed tissues and synovial fluid from patients      |  |
| with rheumatoid arthritis. As such, it is believed to play an important role in the regulation of |  |
| inflammatory diseases. The formation of PLAP can be stimulated by IL-1 beta and TNF-alpha.        |  |
|   |  |

Gene ID: 9373

UniProt: Q9Y263

## **Application Details**

Application Notes: WB 1:300-5000

IHC-P 1:200-400 IHC-F 1:100-500

Restrictions: For Research Use only

## Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Handling Advice:   | Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.               |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |