antibodies

## Datasheet for ABIN704516 anti-Presenilin 2 antibody (AA 51-150)



Overview

Quantity:	100 μL
Target:	Presenilin 2 (PSEN2)
Binding Specificity:	AA 51-150
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Presenilin 2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human presenilin 2/PS-2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	Presenilin 2 (PSEN2)
Alternative Name:	presenilin 2/PS-2 (PSEN2 Products)
Background:	Synonyms: AD 3L, AD 3LP, AD 4, AD 5, AD3L, AD3LP, AD4, AD5, Alzheimer disease 4, Alzheimer

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN704516 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

	disease familial type 4, E5 1, Presenilin2, Presenilin 2, Presenilin 2, PS 2, PS2, PSEN 2, PSNL 2,
	PSNL2, STM 2, STM2.
	Background: Alzheimer's disease (AD) patients with an inherited form of the disease carry
	mutations in the presenilin proteins (PSEN1, PSEN2) or the amyloid precursor protein (APP).
	These disease-linked mutations result in increased production of the longer form of amyloid-
	beta (main component of amyloid deposits found in AD brains). Presenilins are postulated to
	regulate APP processing through their effects on gamma-secretase, an enzyme that cleaves
	APP. Also, it is thought that the presenilins are involved in the cleavage of the Notch receptor,
	such that they either directly regulate gamma-secretase activity or themselves are protease
	enzymes. Two alternative transcripts of PSEN2 have been identified.
Gene ID:	5664

Pathways:

Notch Signaling, EGFR Signaling Pathway, Neurotrophin Signaling Pathway

## Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN704516 | 03/07/2024 | Copyright antibodies-online. All rights reserved.