antibodies -online.com





anti-Presenilin 2 antibody (AA 51-150) (Biotin)



Go to Product page

()	1/0	r\ / I	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	Presenilin 2 (PSEN2)	
Binding Specificity:	AA 51-150	
Reactivity:	Human, Rat, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Presenilin 2 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human presenilin 2/PS-2	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

Target Details

Target:	Presenilin 2 (PSEN2)	
Alternative Name:	presenilin 2/PS-2 (PSEN2 Products)	
Background:	Synonyms: AD 3L, AD 3LP, AD 4, AD 5, AD3L, AD3LP, AD4, AD5, Alzheimer disease 4, Alzheimer	
	disease familial type 4, E5 1, Presenilin2, Presenilin 2, Presenilin 2, PS 2, PS2, PSEN 2, PSNL 2,	

PSNL2.	STM 2	CMT2
FOINLZ.	\sim 1 IVI \sim	. OTTVIZ.

Background: Alzheimer's disease (AD) patients with an inherited form of the disease carry mutations in the presenilin proteins (PSEN1, PSEN2) or the amyloid precursor protein (APP). These disease-linked mutations result in increased production of the longer form of amyloid-beta (main component of amyloid deposits found in AD brains). Presenilins are postulated to regulate APP processing through their effects on gamma-secretase, an enzyme that cleaves APP. Also, it is thought that the presenilins are involved in the cleavage of the Notch receptor, such that they either directly regulate gamma-secretase activity or themselves are protease enzymes. Two alternative transcripts of PSEN2 have been identified.

Gene ID:

5664

Pathways:

Notch Signaling, EGFR Signaling Pathway, Neurotrophin Signaling Pathway

Application Details

Application Notes:

WB 1:300-5000

IHC-P 1:200-400

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months