# antibodies -online.com





## anti-UGT2B4 antibody (AA 111-210)



Go to Product page

$\sim$			
	N/P	r\/I	$\Theta \backslash \Lambda /$

Quantity:	100 μL
Target:	UGT2B4
Binding Specificity:	AA 111-210
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UGT2B4 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human UGT2B4
Isotype:	IgG
Predicted Reactivity:	Human,Sheep
Purification:	Purified by Protein A.

## **Target Details**

Target:	UGT2B4
Alternative Name:	UGT2B4 (UGT2B4 Products)

#### Target Details

Bac	kar	our	nd:
		<b>О О</b> 1	

Synonyms: HLUG25, Hyodeoxycholic acid, UDP glucuronosyltransferase 2 family polypeptide B4, UDP glucuronosyltransferase 2B4 precursor, UDP glucuronyltransferase family 2 beta 4, UDP glycosyltransferase 2 family polypeptide B4, UDP glycosyltransferase, UDPGT, UDPGTh 1, UDPGTh1, UGT2B11, UGT2B4, UD2B4\_HUMAN.

Background: The UDP-Glucuronosyltransferases (UGT) comprise a family of enzymes that detoxify and enhance the urinary excretion of a wide variety of xenobiotic and endogenous substrates by transferring glucuronic acid to sulfhydryl, hydroxyl, aromatic amino, or carboxylic acid groups. They have been subdivided into two families, UGT1 and UGT2, based on the evolutionary divergence of their genes. The enzymes of the UGT1A family play an important role in the metabolism of dietary constituents, phenols, and therapeutic drugs, and also the glucuronidation of bilirubin and iodothyronines. The enzymes of the UGT2B family are involved in the metabolism of bile acids, phenol derivatives, catecholestrogens and steroids. Although it is widely recognized that the liver is the major site of glucuronidation, it is now clear that UGT enzymes are also found in extra-hepatic tissues.

Gene ID:

7363

Pathways:

Steroid Hormone Biosynthesis

#### **Application Details**

Λ	nlination	Matan
ΑD	plication	notes.

WB 1:300-5000

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

## Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months