

[Go to Product page](#)

Datasheet for ABIN706123

**anti-TRIP10 antibody (AA 151-250) (Biotin)**

## Overview

Quantity:	100 µL
Target:	TRIP10
Binding Specificity:	AA 151-250
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRIP10 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Cip4
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Pig,Horse
Purification:	Purified by Protein A.

## Target Details

Target:	TRIP10
Alternative Name:	Cip4 ( <a href="#">TRIP10 Products</a> )

## Target Details

Background:	<p>Synonyms: Cdc42 interacting protein 4, Cdc42 interaction protein 4 long isoform, Cdc42-interacting protein 4, Cip 4, CIP4_HUMAN, DCIP4, hSTP, Protein Felic, Salt tolerant protein, Salt tolerator, STOT, STP, Thyroid hormone receptor interactor 10, Thyroid receptor interacting protein 10, Thyroid receptor-interacting protein 10, TR-interacting protein 10, TRIP 10, TRIP-10, trip10, Truncated Cdc42 interaction protein 4.</p> <p>Background: Required for translocation of GLUT4 to the plasma membrane in response to insulin signaling (By similarity). Required to coordinate membrane tubulation with reorganization of the actin cytoskeleton during endocytosis. Binds to lipids such as phosphatidylinositol 4,5-bisphosphate and phosphatidylserine and promotes membrane invagination and the formation of tubules. Also promotes CDC42-induced actin polymerization by recruiting WASL/N-WASP which in turn activates the Arp2/3 complex. Actin polymerization may promote the fission of membrane tubules to form endocytic vesicles. Required for the formation of podosomes, actin-rich adhesion structures specific to monocyte-derived cells. May be required for the lysosomal retention of FASLG/FASL.</p>
Gene ID:	9322

## Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

## Handling

---

Expiry Date: 12 months