antibodies - online.com







anti-ELAC2 antibody (AA 88-190) (Biotin)



Overview

Quantity:	100 μL
Target:	ELAC2
Binding Specificity:	AA 88-190
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ELAC2 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ELAC2
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	ELAC2
Alternative Name:	ELAC2 (ELAC2 Products)

Target Details

Preservative:

Storage:

Expiry Date:

Precaution of Use:

Storage Comment:

rarget Details	
Background:	Synonyms: elaC homolog 2, ElaC homolog protein 2, ELC2, Heredity prostate cancer protein 2,
	HPC2, putative prostate cancer susceptibility protein HPC2/ELAC2, Ribonuclease Z2, RNase Z2
	tRNA 3 endonuclease 2, tRNase Z long form, tRNase Z2, Zinc phosphodiesterase ELAC protein
	2.
	Background: ELAC2 is a zinc phosphodiesterase which displays some tRNA 3' processing
	endonuclease activity. It is thought to be involved in tRNA maturation, acting by removing a 3'
	trailer from precursor tRNA. ELAC2 also interacts with activated Smad family member 2
	(Smad2) and its nuclear partner forkhead box H1 (also known as FAST-1), and reduced
	expression can suppress transforming growth factor-beta induced growth arrest. Mutations in
	this gene result in an increased risk of prostate cancer. A number of allelic variants have been
	investigated in hereditary prostate cancer.
Gene ID:	60528
Application Details	
Application Notes:	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

50 % Glycerol.

handled by trained staff only.

Store at -20°C for 12 months.

ProClin

-20 °C

12 months