antibodies -online.com





Datasheet for ABIN706333 anti-NCK2 antibody (AA 301-380) (Biotin)

Go to Product page

()	1 /	0	rv	/ 1 /	71	Α.
	1//	\vdash	1 \/	16		1/1/
\sim	v	\sim	1 V	١,	_	v v

Quantity:	100 μL	
Target:	NCK2	
Binding Specificity:	AA 301-380	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NCK2 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NCK2
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NCK2
Alternative Name:	NCK2 (NCK2 Products)

Target Details

Background:	Synonyms: Cytoplasmic protein NCK2, NCK2, GRB4, Growth factor receptor bound protein 4,
	NCK adaptor protein 2, Nck-2, NCK2, NCK2_HUMAN, NCKbeta, Non-catalytic region of tyrosine
	kinase adaptor protein 2, Noncatalytic region of tyrosine kinase beta, SH2/SH3 adaptor protein
	NCK-beta, SH2/SH3 adaptor protein NCK-beta.
	Background: This gene encodes a member of the NCK family of adaptor proteins. The protein
	contains three SH3 domains and one SH2 domain. The protein has no known catalytic function
	but has been shown to bind and recruit various proteins involved in the regulation of receptor
	protein tyrosine kinases. It is through these regulatory activities that this protein is believed to
	be involved in cytoskeletal reorganization. Alternate transcriptional splice variants, encoding
	different isoforms, have been characterized. [provided by RefSeq].
Gene ID:	8440
Pathways:	EGFR Signaling Pathway, Regulation of Actin Filament Polymerization
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months