# antibodies -online.com







# anti-HNRNPK antibody (AA 201-300)





#### Overview

Quantity:	100 μL
Target:	HNRNPK
Binding Specificity:	AA 201-300
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HNRNPK antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human hnRNP K
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Pig,Chicken,Rabbit
Purification:	Purified by Protein A.

# **Target Details**

|--|


# **Target Details**

Alternative Name:	hnRNP K (HNRNPK Products)
Background:	Synonyms: CSBP, TUNP, HNRPK, Heterogeneous nuclear ribonucleoprotein K, hnRNP K,
	Transformation up-regulated nuclear protein, HNRNPK
	Background: One of the major pre-mRNA-binding proteins. Binds tenaciously to poly(C)
	sequences. Likely to play a role in the nuclear metabolism of hnRNAs, particularly for pre-
	mRNAs that contain cytidine-rich sequences. Can also bind poly(C) single-stranded DNA. Plays
	an important role in p53/TP53 response to DNA damage, acting at the level of both
	transcription activation and repression. When sumoylated, acts as a transcriptional coactivato
	of p53/TP53, playing a role in p21/CDKN1A and 14-3-3 sigma/SFN induction (By similarity). As
	far as transcription repression is concerned, acts by interacting with long intergenic RNA p21
	(lincRNA-p21), a non-coding RNA induced by p53/TP53. This interaction is necessary for the
	induction of apoptosis, but not cell cycle arrest.
Gene ID:	3190
UniProt:	P61978
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.

## Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

# **Images**



## **Western Blotting**

**Image 1.** Lane 1:A549 cell lysates probed with Rabbit Anti-hnRNP K Polyclonal Antibody, Unconjugated (ABIN706571) at 1:300 overnight at 4 °C. Followed by conjugation to secondary antibody at 1:5000 for 90 min at 37 °C.