

Datasheet for ABIN7072417

Recombinant anti-CD40 Ligand antibody



[Go to Product page](#)

1 Image

Overview

Quantity:	200 µg
Target:	CD40 Ligand (CD40LG)
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Chimeric
Conjugate:	This CD40 Ligand antibody is un-conjugated
Application:	ELISA

Product Details

Purpose:	Anti-CD40L [AT161-10], Rabbit IgG, kappa
Immunogen:	This antibody was raised by immunising mice with Chinese hamster ovary (CHO) cells expressing human CD40L.
Clone:	AT161-10
Isotype:	IgG kappa
Specificity:	This antibody is specific for human CD40L, and has been screened for specificity using human CD40L-rat CD40 fusion protein
Characteristics:	Original Species of Ab: Mouse Original Format of Ab: IgG1
Purification:	Protein A affinity purified

Target Details


Target:	CD40 Ligand (CD40LG)
Alternative Name:	CD40L (CD40LG Products)
Background:	CD154, CD40 ligand, CD-40L, CD-40 Ligand, CD40-L, CD-40-L
Pathways:	NF-kappaB Signaling , Production of Molecular Mediator of Immune Response , Cancer Immune Checkpoints

Application Details

Application Notes:	This antibody has been used in an ELISA analysis to quantify serum levels of CD40L in samples from Alzheimer's patients (direct communication with originator, 2017).
Comment:	This chimeric rabbit antibody was made using the variable domain sequences of the original Mouse IgG1 format, for improved compatibility with existing reagents, assays and techniques.
Restrictions:	For Research Use only

Handling

Concentration:	1 mg/mL
Buffer:	PBS with 0.02 % Proclin 300.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.



Immunofluorescence

Image 1. Immunofluorescence staining of human peripheral blood monocytes using anti-CD40L antibody AT161-10. Immunofluorescence analysis of paraformaldehyde fixed human peripheral blood monocytes immobilized on Shi-fix™ cover-slips and stained with the chimeric rabbit IgG version of AT161-10 (ABIN7072417) at 10 µg/mL followed by Alexa Fluor® 488 secondary antibody (2 µg/mL), showing membrane staining. The nuclear stain is DAPI (blue). Panels show from left-right, top-bottom ABIN7072417, DAPI, merged channels and an isotype control. The isotype control was stained with anti-Fluorescein antibody (ABIN5668035) followed by Alexa Fluor® 488 secondary antibody.