antibodies - online.com





anti-CYP46A1 antibody (AA 51-150)



Image



Overview

Quantity:	100 μL
Target:	CYP46A1
Binding Specificity:	AA 51-150
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CYP46A1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CYP46
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Cow,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target Details

Alternative Name:	CYP46 (CYP46A1 Products)
Background:	Synonyms: Cholesterol 24-hydroxylase, CP46, CYP46, cytochrome P450 family 46,
	CP46A_HUMAN.
	Background: The Cyp46 gene encodes a member of the cytochrome P450 superfamily of
	enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions
	involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This
	endoplasmic reticulum protein is expressed exclusively in the brain, where it converts
	cholesterol to 24S-hydroxycholesterol by adding a hydroxyl group to cholesterol, producing a
	product that is more soluble than cholesterol and able to be exported from the brain.Cyp46 is
	also known as 24S-cholesterol hydroxylase.
Gene ID:	10858
Application Details	

Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date:

12 months

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Paraformaldehyde-fixed, paraffin embedded Mouse brain, Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min, Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes, Blocking buffer (normal goat serum) at 37°C for 30min, Antibody incubation with CYP46 Polyclonal Antibody, Unconjugated at 1:400 overnight at 4°C, followed by a conjugated secondary antibody (sp-0023) for 20 minutes and DAB staining.