

Datasheet for ABIN7091120 anti-FOXO1 antibody (pSer256) (PE)



Overview

Quantity:	100 μL
Target:	FOX01
Binding Specificity:	pSer256
Reactivity:	Human, Mouse, Rat, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FOXO1 antibody is conjugated to PE
Application:	Western Blotting (WB), Flow Cytometry (FACS)
Product Details	
Immunogen:	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human FOXO1 around the
lmmunogen:	KLH conjugated synthetic phosphopeptide derived from human FOXO1 around the phosphorylation site of Ser256 [AA(p-S)MD]
Immunogen: Isotype:	
	phosphorylation site of Ser256 [AA(p-S)MD]
Isotype:	phosphorylation site of Ser256 [AA(p-S)MD]
Isotype: Cross-Reactivity:	phosphorylation site of Ser256 [AA(p-S)MD] IgG Cow, Human, Mouse, Rat
Isotype: Cross-Reactivity: Predicted Reactivity:	phosphorylation site of Ser256 [AA(p-S)MD] IgG Cow, Human, Mouse, Rat Dog,Cow,Pig,Horse,Chicken
Isotype: Cross-Reactivity: Predicted Reactivity: Purification:	phosphorylation site of Ser256 [AA(p-S)MD] IgG Cow, Human, Mouse, Rat Dog,Cow,Pig,Horse,Chicken

Target Details

Target Details	
Background:	Synonyms: FOXO1A phospho S256, p-FOXO1A phospho S256, FKHRSer256, Forkhead box
	protein O1, Afxh, Al876417, FKHR, Fkhr1, Foxo1a, Forkhead, FKH 1, FKH1, FKH1, FKHR, FKHR,
	Forkhead Drosophila homolog 1 rhabdomyosarcoma, Forkhead Drosophila homolog 1
	rhabdomyosarcoma, Forkhead box 01, Forkhead box protein 01, Forkhead box protein 01A,
	Forkhead in rhabdomyosarcoma, Forkhead, Drosophila, homolog of, in rhabdomyosarcoma,
	FOXO1, FOXO1_HUMAN, FOXO1A.
	Background: This gene belongs to the forkhead family of transcription factors which are
	characterized by a distinct forkhead domain. The specific function of this gene has not yet been
	determined, however, it may play a role in myogenic growth and differentiation. Translocation
	of this gene with PAX3 has been associated with alveolar rhabdomyosarcoma. [provided by
	RefSeq].
Gene ID:	2308
Pathways:	PI3K-Akt Signaling, Cell Division Cycle, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling
	Pathway, Neurotrophin Signaling Pathway, Carbohydrate Homeostasis, Chromatin Binding,
	Regulation of Carbohydrate Metabolic Process, CXCR4-mediated Signaling Events, BCR
	Signaling
Application Details	
Application Notes:	FCM 1:20-100
Restrictions:	For Research Use only
Handling	
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.

Preservative:

ProClin

Precaution of Use:

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be

handled by trained staff only.

Storage:

-20 °C

Storage Comment:

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.