antibodies .- online.com





Go to Product page

Datasheet for ABIN709151

anti-HADHB antibody (AA 231-330)

Overview 100 μL Quantity: Target: **HADHB** Binding Specificity: AA 231-330 Reactivity: Human, Mouse Host: Rabbit Clonality: Polyclonal Conjugate: This HADHB antibody is un-conjugated Application: Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffinembedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human HADHB
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target Details

Alternative Name:	HADHB (HADHB Products)
Background:	Synonyms: 2 enoyl Coenzyme A (CoA) hydratase beta subunit, 3 ketoacyl Coenzyme A (CoA)
	thiolase of mitochondrial trifunctional protein beta subunit, 3 ketoacyl Coenzyme A thiolase, 3
	ketoacyl Coenzyme A thiolase of mitochondrial trifunctional protein beta subunit, Acetyl CoA
	acyltransferase, Beta ketothiolase, ECHB, HADH, Hydroxyacyl Coenzyme A (CoA)
	dehydrogenase beta subunit, Hydroxyacyl Coenzyme A dehydrogenase, Hydroxyacyl Coenzyme
	A dehydrogenase beta subunit, Hydroxyacyl Coenzyme A dehydrogenase/3 ketoacyl Coenzyme
	A thiolase/enoyl Coenzyme A hydratase (trifunctional protein) beta subunit, Mitochondrial
	trifunctional enzyme beta subunit, Mitochondrial trifunctional protein beta subunit, MTPB, TP
	beta. TPbeta, Trifunctional enzyme subunit beta, Trifunctional enzyme subunit beta
	mitochondrial, Trifunctional protein, ECHB_HUMAN.
	Background: The HADHB gene encodes the beta subunit of the mitochondrial trifunctional
	protein, which catalyzes the last three steps of mitochondrial beta-oxidation of long chain fatty
	acids. The mitochondrial membrane-bound heterocomplex is composed of four alpha and four
	beta subunits, with the beta subunit catalyzing the 3-ketoacyl-CoA thiolase activity. Mutations is
	this gene result in trifunctional protein deficiency. The encoded protein can also bind RNA and
	decreases the stability of some mRNAs. The genes of the alpha and beta subunits of the
	mitochondrial trifunctional protein are located adjacent to each other in the human genome in
	head-to-head orientation. Alternatively spliced transcript variants have been found, however,
	their full-length nature is not known.
Gene ID:	151556
UniProt:	P55084
Pathways:	Monocarboxylic Acid Catabolic Process
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months