antibodies .- online.com





anti-PKD2 antibody (AA 41-140) (Biotin)



Go to Product page

()	11/	NI	/1/	≥W
	/ // 	۰۱ ا	// I E	- ¹ \/ \/

Quantity:	100 μL
Target:	PKD2
Binding Specificity:	AA 41-140
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PKD2 antibody is conjugated to Biotin
Application:	ELISA, Western Blotting (WB)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Polycystin 2
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Cow,Pig,Horse,Rabbit,Zebrafish
Purification:	Purified by Protein A.

Target Details

Target:	PKD2
Alternative Name:	Polycystin 2 (PKD2 Products)
Background: Synonyms: polycystic kidney disease 2, TRPP2, APKD2, C030034P18RIK, Mi	

MGC138468, PC2, PKD2 (includes EG:5311), PKD4, POLYCISTIN-2, POLYCYSTIN 2,
RGD1559992, TRPP2, Polycystic kidney disease 2 protein homolog, PC2.
Background: Involved in fluid-flow mechanosensation by the primary cilium in renal epithelium.
PKD1 and PKD2 may function through a common signaling pathway that is necessary for
normal tubulogenesis. Acts as a regulator of cilium length, together with PKD1. The dynamic
control of cilium length is essential in the regulation of mechanotransductive signaling. The
cilium length response creates a negative feedback loop whereby fluid shear-mediated
deflection of the primary cilium, which decreases intracellular cAMP, leads to cilium shortening
and thus decreases flow-induced signaling. Functions as a calcium permeable cation channel.
5311

Gene ID:	5311	
UniProt:	Q13563	
Pathways:	cAMP Metabolic Process, Maintenance of Protein Location, Negative Regulation of Transporter	
	Activity	

Application Details

Application Notes:	WB 1:300-5000
Restrictions:	For Research Use only
Handling	
Concentration:	1 μg/μL

Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	