antibodies .- online.com







anti-HMGCL antibody (AA 41-140) (Biotin)



Overview

Quantity:	100 μL
Target:	HMGCL
Binding Specificity:	AA 41-140
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HMGCL antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HMGCL
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	HMGCL
Alternative Name:	HMGCL (HMGCL Products)

Target Details

Background:	Synonyms: 3 hydroxy 3 methylglutaryl CoA lyase, 3 hydroxy 3 methylglutaryl Coenzyme A lyase,
	3 hydroxymethyl 3 methylglutaryl Coenzyme A lyase hydroxymethylglutaricaciduria, 3
	hydroxymethyl 3 methylglutaryl Coenzyme A lyase, 3-hydroxy-3-methylglutarate-CoA lyase, HL,
	HMG CoA lyase, HMG CoA Lyase Deficiency, HMG-CoA lyase, HMGCL, HMGCL_HUMAN,
	Hydroxymethylglutaricaciduria, Hydroxymethylglutaryl CoA lyase, Hydroxymethylglutaryl CoA
	lyase mitochondrial, Hydroxymethylglutaryl-CoA lyase, mitochondrial, MS725,
	OTTHUMP0000044830.
	Background: Hydroxymethylglutaryl-CoA lyase (HMGCL) is found in fibroblasts, liver and
	lymphoblasts. It has a role in ketogenesis and leucine catabolism. Defects in HMGCL are the
	cause of 3-hydroxy-3-methylglutaryl-CoA lyase deficiency (hydroxymethylglutaricaciduria), an
	autosomal recessive disease which can lead to hypoglycemia and coma.
Gene ID:	2835
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months