

Datasheet for ABIN7101704
anti-U2AF2 antibody



[Go to Product page](#)

4 Images

Overview

Quantity:	20 µL
Target:	U2AF2 (U2AF59)
Reactivity:	Human
Host:	Rabbit
Clonality:	Monoclonal
Conjugate:	This U2AF2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)

Product Details

Immunogen:	A synthesized peptide derived from human U2AF2/U2AF2/U2AF65
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Monoclonal Antibodies
Purification:	Affinity purification

Target Details

Target:	U2AF2 (U2AF59)
Alternative Name:	U2AF2 (U2AF59 Products)
Background:	U2 auxiliary factor (U2AF), comprised of a large and a small subunit, is a non-snRNP protein required for the binding of U2 snRNP to the pre-mRNA branch site. This gene encodes the U2AF

Target Details

large subunit which contains a sequence-specific RNA-binding region with 3 RNA recognition motifs and an Arg/Ser-rich domain necessary for splicing. The large subunit binds to the polypyrimidine tract of introns early during spliceosome assembly. Multiple transcript variants have been detected for this gene, but the full-length natures of only two have been determined to date. [provided by RefSeq, Jul 2008],U2AF65,Epigenetics & Nuclear Signaling,RNA Binding,U2AF2

Molecular Weight: 54 kDa

Gene ID: 11338

UniProt: [P26368](#)

Application Details

Application Notes: WB,1:500 - 1:2000,IHC,1:50 - 1:200,IP,1:50 - 1:200

Restrictions: For Research Use only

Handling


Buffer: PBS with 0.02 % sodium azide,0.05 % BSA,50 % glycerol, pH 7.3.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C

Storage Comment: Store at -20°C. Avoid freeze / thaw cycles.




Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human colon using U2/U2/U2 Rabbit mAb (ABIN1682786, ABIN3019042, ABIN3019043 and ABIN7101704) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Western Blotting

Image 2. Western blot analysis of extracts of various cell lines, using U2/U2/U2 Rabbit mAb (ABIN1682786, ABIN3019042, ABIN3019043 and ABIN7101704) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.



Western Blotting

Image 3. Western blot analysis of extracts of various cell lines, using U2/U2/U2 Rabbit mAb (ABIN1682786, ABIN3019042, ABIN3019043 and ABIN7101704) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN7101704.