

Datasheet for ABIN710283

anti-MAPK7 antibody (pSer731, pThr733) (FITC)[Go to Product page](#)**1** Image

Overview

Quantity:	100 µL
Target:	MAPK7
Binding Specificity:	pSer731, pThr733
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAPK7 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human ERK5 around the phosphorylation sites of Ser731+Thr733
Isotype:	IgG
Specificity:	Recognizes Ser721+Thr723 in mice and rats
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	MAPK7
Alternative Name:	ERK5+ (MAPK7 Products)
Background:	<p>Synonyms: BMK1, ERK4, ERK5, PRKM7, Mitogen-activated protein kinase 7, MAP kinase 7, MAPK 7, Big MAP kinase 1, BMK-1, Extracellular signal-regulated kinase 5, ERK-5, MAPK7</p> <p>Background: Plays a role in various cellular processes such as proliferation, differentiation and cell survival. The upstream activator of MAPK7 is the MAPK kinase MAP2K5. Upon activation, it translocates to the nucleus and phosphorylates various downstream targets including MEF2C. EGF activates MAPK7 through a Ras-independent and MAP2K5-dependent pathway. May have a role in muscle cell differentiation. May be important for endothelial function and maintenance of blood vessel integrity. MAP2K5 and MAPK7 interact specifically with one another and not with MEK1/ERK1 or MEK2/ERK2 pathways. Phosphorylates SGK1 at Ser-78 and this is required for growth factor-induced cell cycle progression. Involved in the regulation of p53/TP53 by disrupting the PML-MDM2 interaction.</p>
Gene ID:	5598
UniProt:	Q13164
Pathways:	MAPK Signaling , Neurotrophin Signaling Pathway , Activation of Innate immune Response , cAMP Metabolic Process , Toll-Like Receptors Cascades , Negative Regulation of intrinsic apoptotic Signaling

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------


Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months

Images



Immunofluorescence (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin embedded human kidney labeled with Rabbit Anti-ERK5 (Ser731+Thr733) Polyclonal Antibody, FITC Conjugated (ABIN710283) at 1:200 followed by conjugation to the secondary antibody and DAB staining