

Datasheet for ABIN710688

anti-NFKBIA antibody (pTyr42) (FITC)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	NFKBIA
Binding Specificity:	pTyr42
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NFKBIA antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Flow Cytometry (FACS)

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human IKB alpha around the phosphorylation site of Tyr42
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Cow, Sheep, Pig, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NFKBIA
---------	--------

Target Details

Alternative Name:	NFKBIA (NFKBIA Products)
Background:	Synonyms: NFKBIATyr42, NFKBIaphospho Y42, IKB-alphaTyr42, IKB alpha, Inhibitor of KB alpha, I kappa B alpha, IKappaBalpaha, IkappaBalpaha, IKBA, IKBalpha, MAD 3, MAD3, Major histocompatibility complex enhancer binding protein MAD3, NF kappa B inhibitor alpha, NFKBI, NFKBIA, Nuclear factor of kappa light chain gene enhancer in B cells, Nuclear factor of kappa light polypeptide gene enhancer in B cells inhibitor alpha. Background: This gene encodes a member of the NF-kappa-B inhibitor family, which contain multiple ankrin repeat domains. The encoded protein interacts with REL dimers to inhibit NF-kappa-B/REL complexes which are involved in inflammatory responses. The encoded protein moves between the cytoplasm and the nucleus via a nuclear localization signal and CRM1-mediated nuclear export. Mutations in this gene have been found in ectodermal dysplasia anhidrotic with T-cell immunodeficiency autosomal dominant disease. [provided by RefSeq, Aug 2011].
Gene ID:	4792
Pathways:	NF-kappaB Signaling , TCR Signaling , TLR Signaling , Fc-epsilon Receptor Signaling Pathway , Activation of Innate immune Response , Cellular Response to Molecule of Bacterial Origin , Maintenance of Protein Location , Hepatitis C , Protein targeting to Nucleus , Toll-Like Receptors Cascades , BCR Signaling

Application Details

Application Notes:	FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months