antibodies -online.com





anti-NDEL1 antibody (pSer231) (Biotin)



()	11/	IN	/ie	A .
	/ // 	۱ ات	/ (−	' \/\/

Overview		
Quantity:	100 μL	
Target:	NDEL1	
Binding Specificity:	pSer231	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NDEL1 antibody is conjugated to Biotin	
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	
Product Details		
lmmunogen:	KLH conjugated synthetic phosphopeptide derived from human NDEL1 around the phosphorylation site of Ser231	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human, Mouse, Dog, Cow, Pig, Horse, Chicken, Rabbit	
Purification:	Purified by Protein A.	
Target Details		
Target:	NDEL1	

Target Details

Alternative Name:	NDEL1 (NDEL1 Products)	
Background:	Synonyms: NDEL1phospho S231, Nudelphospho S231, A. nidulans, DKFZp451M0318,	
	ENDOOLIGOPEPTIDASE A, EOPA, MITAP 1, MITAP1, Mitosin associated protein 1, Mitosin	
	associated protein MITAP1, Mitosin-associated protein 1, Ndel 1, Ndel1, NDEL1_HUMAN,	
	Nuclear distribution gene E like homolog 1, Nuclear distribution protein nudE like 1, Nuclear	
	distribution protein nudE-like 1, NUDE like protein, NudE nuclear distribution gene E homolog	
	like 1 A. nidulans, NudE nuclear distribution gene E homolog like 1, NUDEL, Protein Nudel.	
	Background: Nudel is important for normal cortical development. It is invovled in microtubule	
	organization, nuclear translocation, and neuronal positioning in concert with various other	
	factors (including Lis1, Pafah1b1, Pahfah1b2, dynein, dynorphin A and cdk5). Western blot	
	analysis of mouse tissues shows abundant expression of Nudel in brain and testis, and much	
	lower expression in heart, liver, kidney, and skeletal muscle. In fractionated rat brain, Nudel and	
	Lis1 are both found in fractions enriched for postsynaptic density proteins. Immunostaining of	
	embryonic day 18 mouse brain sections revealed staining of migrating neurons and	
	thalamocortical axons of the intermediate zone of the developing cerebral cortex, as well as	
	several other developing brain regions. The deduced protein contains 345 amino acids and has	
	a calculated molecular mass of 38.4 kDa. It has a coiled coil motif (residues 19 to 201),	
	followed by several potential phosphorylation sites for casein kinase II, protein kinase C or	
	CDK5. Nudel shares about 50 % identity with mouse and human NUDE proteins.	
Gene ID:	81565	
Pathways:	Regulation of Cell Size	
Application Details		
Application Notes:	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 an	
	50 % Glycerol.	

Handling

Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	