

Datasheet for ABIN711348

anti-PTPN7 antibody (pSer93) (FITC)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	PTPN7
Binding Specificity:	pSer93
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PTPN7 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human PTPN7 around the phosphorylation site of Ser93
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow
Purification:	Purified by Protein A.

Target Details

Target:	PTPN7
Alternative Name:	PTPN7 (PTPN7 Products)

Target Details

Background:	Synonyms: PTPN7phospho S93, BPTP 4, BPTP4, Dual specificity phosphatase 1, Hematopoietic protein tyrosine phosphatase, LC PTP, LCPTP, LPTP, Protein tyrosine phosphatase non receptor type stress induced, Protein tyrosine phosphatase nonreceptor type stress induced, Protein tyrosine phosphatase LC PTP, Protein tyrosine phosphatase non receptor type 7, PTPN 7, PTPNI, Tyrosine protein phosphatase non receptor type 7, PTN7_HUMAN. Background: The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This gene is preferentially expressed in a variety of hematopoietic cells, and is an early response gene in lymphokine stimulated cells. The non-catalytic N-terminus of this PTP can interact with MAP kinases and suppress the MAP kinase activities. This PTP was shown to be involved in the regulation of T cell antigen receptor (TCR) signaling, which was thought to function through dephosphorylating the molecules related to MAP kinase pathway. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq].
-------------	--

Gene ID:	5778
----------	------

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months