

Datasheet for ABIN711380

anti-PKC zeta antibody (pThr410) (HRP)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	PKC zeta (PRKCZ)
Binding Specificity:	pThr410
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PKC zeta antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human PRKCZ around the phosphorylation site of Thr410
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Horse, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	PKC zeta (PRKCZ)
Alternative Name:	PRKCZ (PRKCZ Products)

Target Details

Background:	Synonyms: PRKCZphospho T410, PRKCZphospho Thr410, p-PRKCZT410, rotein Kinase C, AAG6, Aging associated gene 6, MGC129900, MGC129901, MGC41878, MGC57564, PKC B, PKC beta, PKC zeta, PKC2, PKCB, PKCC, PKCD, PKCE, PKCG, PRKACA, PRKC A, PRKC G, PRKCA, PRKCB, PRKCB1, PRKCB2, PRKCD, PRKCE, PRKCG, PRKCZ , Protein kinase C alpha, Protein kinase C alpha type, Protein Kinase C, Protein kinase C beta 1, Protein kinase C beta 1 polypeptide, Protein kinase C beta, Protein kinase C beta type, Protein kinase C delta, Protein kinase C epsilon, Protein kinase C gamma, Protein kinase C gamma type, Protein kinase C zeta, SCA14, Spinocerebellar ataxia 14. Background: Protein kinase C (PKC) zeta is a member of the PKC family of serine/threonine kinases which are involved in a variety of cellular processes such as proliferation, differentiation and secretion. Unlike the classical PKC isoenzymes which are calcium-dependent, PKC zeta exhibits a kinase activity which is independent of calcium and diacylglycerol but not of phosphatidylserine. Furthermore, it is insensitive to typical PKC inhibitors and cannot be activated by phorbol ester. Unlike the classical PKC isoenzymes, it has only a single zinc finger module. These structural and biochemical properties indicate that the zeta subspecies is related to, but distinct from other isoenzymes of PKC. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008].
-------------	--

Gene ID:	5590
Pathways:	NF-kappaB Signaling , RTK Signaling , Myometrial Relaxation and Contraction , Regulation of Leukocyte Mediated Immunity , Positive Regulation of Immune Effector Process , Synaptic Membrane , Production of Molecular Mediator of Immune Response , CXCR4-mediated Signaling Events , Thromboxane A2 Receptor Signaling

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
--------------------	---

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and

Handling

	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months