antibodies -online.com





anti-EDF1 antibody



| \sim | |
|--------|-------|
| ()\/\ | rview |
| OVC | VICVV |

| Quantity: | 100 μg |
|-----------------|--|
| Target: | EDF1 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This EDF1 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), ELISA |
| Product Details | |
| Immunogen: | endothelial differentiation-related factor 1 |

Purification:

Immunogen affinity purified

IgG

Purity:

Isotype:

≥95 % as determined by SDS-PAGE

Target Details

| Target: | EDF1 |
|-------------------|--|
| Alternative Name: | EDF1 (EDF1 Products) |
| Background: | Synonyms:CFAP280 Background:This gene encodes a protein that may regulate endothelial cell differentiation, lipid metabolism, and hormone-induced cardiomyocyte hypertrophy. The |
| | encoded protein has also been found to act as a transcriptional coactivator by interconnecting |
| | the general transcription factor TATA element-binding protein (TBP) and gene-specific |

Target Details

| | activators. Alternate splicing results in multiple transcript variants. |
|-------------------|---|
| Molecular Weight: | 16 kDa |
| Gene ID: | 8721 |
| UniProt: | 060869 |

Application Details

| Application Notes: | WB: 1:500 - 1:2000, IHC: 1:50 - 1:200 | |
|--------------------|---------------------------------------|--|
| Restrictions: | For Research Use only | |

Handling

| Format: | Liquid |
|--------------------|--|
| Buffer: | PBS with 0.02 % sodium azide and 50 % glycerol pH 7.3, |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | -20°C for 12 months (Avoid repeated freeze / thaw cycles.) |
| Expiry Date: | 12 months |