# antibodies -online.com





## anti-EIF3B antibody



Go to Product page

| $\sim$ |     |     |     |
|--------|-----|-----|-----|
|        | N/P | r\/ | i⊢₩ |

| Quantity:    | 100 μg                               |
|--------------|--------------------------------------|
| Target:      | EIF3B                                |
| Reactivity:  | Human, Mouse, Rat                    |
| Host:        | Rabbit                               |
| Clonality:   | Polyclonal                           |
| Conjugate:   | This EIF3B antibody is un-conjugated |
| Application: | ELISA                                |

### **Product Details**

| Immunogen:    | eukaryotic translation initiation factor 3, subunit B |  |
|---------------|---|--|
| Isotype:      | IgG   |  |
| Purification: | Immunogen affinity purified                           |  |
| Purity:       | ≥95 % as determined by SDS-PAGE                       |  |

## **Target Details**

| Target:           | EIF3B   |
|-------------------|---|
| Alternative Name: | EIF3B (EIF3B Products)  |
| Background:       | Synonyms:EIF3S9 Background:Component of the eukaryotic translation initiation factor 3(eIF-       |
|                   | 3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 |
|                   | complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-   |
|                   | 2:GTP:methionyl-tRNAi and eIF-5 to form the 43S pre-initiation complex(43S PIC). The eIF-3        |

#### **Target Details**

complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. (Microbial infection) In case of FCV infection, plays a role in the ribosomal termination-reinitiation event leading to the translation of VP2(PubMed:18056426).

Gene ID: 8662

UniProt: P55884

Pathways: Ribonucleoprotein Complex Subunit Organization

## **Application Details**

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

## Handling

| Format:            | Liquid   |  |
|--------------------|--|--|
| Buffer:            | PBS with 0.02 % sodium azide and 50 % glycerol pH 7.3,   |  |
| Preservative:      | Sodium azide   |  |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |  |
| Storage:           | -20 °C   |  |
| Storage Comment:   | -20°C for 12 months (Avoid repeated freeze / thaw cycles.)   |  |
| Expiry Date:       | 12 months  |  |