

Datasheet for ABIN7114007 **anti-EXOSC1 antibody**



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	EXOSC1
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EXOSC1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	exosome component 1
Isotype:	IgG
Purification:	Immunogen affinity purified
Purity:	≥95 % as determined by SDS-PAGE

Target Details

Target:	EXOSC1
Alternative Name:	EXOSC1 (EXOSC1 Products)
Background:	Synonyms: CSL4 Background: Non-catalytic component of the RNA exosome complex which has 3'→5' exoribonuclease activity and participates in a multitude of cellular RNA processing and degradation events. In the nucleus, the RNA exosome complex is involved in proper maturation of stable RNA species such as rRNA, snRNA and snoRNA, in the elimination of RNA

Target Details

processing by-products and non-coding 'pervasive' transcripts, such as antisense RNA species and promoter-upstream transcripts(PROMPTs), and of mRNAs with processing defects, thereby limiting or excluding their export to the cytoplasm. The RNA exosome may be involved in Ig class switch recombination(CSR) and/or Ig variable region somatic hypermutation(SHM) by targeting AICDA deamination activity to transcribed dsDNA substrates. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements(AREs) within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems to be involved in degradation of histone mRNA. The catalytic inactive RNA exosome core complex of 9 subunits(Exo-9) is proposed to play a pivotal role in the binding and presentation of RNA for ribonucleolysis, and to serve as a scaffold for the association with catalytic subunits and accessory proteins or complexes. EXOSC1 as peripheral part of the Exo-9 complex stabilizes the hexameric ring of RNase PH-domain subunits through contacts with EXOSC6 and EXOSC8.

Molecular Weight:	21 kDa
Gene ID:	51013
UniProt:	Q9Y3B2

Application Details

Application Notes:	WB: 1:500-1:2000, IHC: 1:20-1:200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	PBS with 0.02 % sodium azide and 50 % glycerol pH 7.3,
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	-20°C for 12 months (Avoid repeated freeze / thaw cycles.)
Expiry Date:	12 months