antibodies .- online.com





anti-EXOSC5 antibody



()	ve	K\ /		A .
	\cup	1 V/	-	V۷

Quantity:	100 μg
Target:	EXOSC5
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EXOSC5 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	exosome component 5
Isotype:	IgG
Purification:	Immunogen affinity purified
Purity:	≥95 % as determined by SDS-PAGE

Target Details

Target:	EXOSC5	
Alternative Name:	EXOSC5 (EXOSC5 Products)	
Background:	Synonyms:CML28, RRP46 Background:Non-catalytic component of the RNA exosome complex	
	which has 3'->5' exoribonuclease activity and participates in a multitude of cellular RNA	
	processing and degradation events. In the nucleus, the RNA exosome complex is involved in	
	proper maturation of stable RNA species such as rRNA, snRNA and snoRNA, in the elimination	

of RNA processing by-products and non-coding 'pervasive' transcripts, such as antisense RNA species and promoter-upstream transcripts(PROMPTs), and of mRNAs with processing defects, thereby limiting or excluding their export to the cytoplasm. The RNA exosome may be involved in Ig class switch recombination(CSR) and/or Ig variable region somatic hypermutation(SHM) by targeting AICDA deamination activity to transcribed dsDNA substrates. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements(AREs) within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems to be involved in degradation of histone mRNA. The catalytic inactive RNA exosome core complex of 9 subunits(Exo-9) is proposed to play a pivotal role in the binding and presentation of RNA for ribonucleolysis, and to serve as a scaffold for the association with catalytic subunits and accessory proteins or complexes.

Molecular Weight:	28-30 kDa
Gene ID:	56915
UniProt:	Q9NQT4
Pathways:	SARS-CoV-2 Protein Interactome

Application Details

Application Notes:	WB: 1:500-1:2000, IHC: 1:20-1:200	
Restrictions:	For Research Use only	

Handling

Format:	Liquid
Buffer:	PBS with 0.02 % sodium azide and 50 % glycerol pH 7.3,
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	-20°C for 12 months (Avoid repeated freeze / thaw cycles.)
Expiry Date:	12 months