

Datasheet for ABIN711635 anti-SHP1 antibody (pTyr536) (HRP)



\sim				
()\	ve	r\/		Λ/
\cup	$V \subset$	ı vı	\Box	٧V

100 μL
SHP1 (PTPN6)
pTyr536
Human, Mouse
Rabbit
Polyclonal
This SHP1 antibody is conjugated to HRP
Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
KLH conjugated synthetic phosphopeptide derived from human PTPN6 around the phosphorylation site of Tyr536 [SE(p-Y)GN]
IgG
Human, Mouse
Dog,Horse
g)· · - ·
Purified by Protein A.

Target Details

Alternative Name:	PTPN6 (PTPN6 Products)		
Background:	Synonyms: SHP-1Tyr536, PTPN6phospho Y536, SHP1phospho Y536, 70 kda SHP1L protein,		
	HCP, HCPH, Hematopoietic cell phosphatase, Hematopoietic cell protein tyrosine phosphatase		
	Hematopoietic cell protein-tyrosine phosphatase, HPTP1C, Protein tyrosine phosphatase 1C,		
	Protein tyrosine phosphatase non receptor type 6, Protein tyrosine phosphatase SHP1, PTP1C,		
	PTPN6, SH PTP 1, SHP1, SHP1L, PTN6_HUMAN.		
	Background: SHP1 is a member of the non receptor protein tyrosine phosphatase subfamily.		
	The PTP family comprises of at least 37 proteins, characterized by a catalytic phosphatase		
	domain of approximately 240 amino acids, and includes both transmembrane and cytosolic		
	enzymes. PTP1B is cytosolic. The PTPs have high substrate specificity for phosphotyrosyl		
	proteins, at the primary sequence level sharing little similarity with the protein serine		
	phosphatases, protein threonine phosphatases, or the acid and alkaline phosphatases. SHP1 is		
	implicated in the control of tyrosine kinase signalling pathways in cellular proliferation, with a		
	potential role in cancer.		
Gene ID:	5777		
UniProt:	P29350		
Pathways:	JAK-STAT Signaling, TCR Signaling, TLR Signaling, Nuclear Receptor Transcription Pathway,		
	Positive Regulation of Peptide Hormone Secretion, Steroid Hormone Mediated Signaling		
	Pathway, Response to Growth Hormone Stimulus, Regulation of Leukocyte Mediated Immunity		
	CXCR4-mediated Signaling Events, Signaling Events mediated by VEGFR1 and VEGFR2, BCR		
	Signaling		
Application Details			
Application Notes:	WB 1:300-5000		
	IHC-P 1:200-400		
	IHC-F 1:100-500		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		

Handling

	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months