antibodies -online.com





anti-SMARCD1 antibody



Go to Product page

\sim			
	N/F	ا/\r14	$\triangle W$

Quantity:	100 μg
Target:	SMARCD1
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SMARCD1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP)

Product Details

Immunogen:	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily d, member 1
Isotype:	IgG
Purification:	Immunogen affinity purified
Purity:	≥95 % as determined by SDS-PAGE

Target Details

Target:	SMARCD1	
Alternative Name:	SMARCD1 (SMARCD1 Products)	
Background:	Synonyms:BAF60A Background:Involved in chromatin remodeling. Belongs to the neural	
	progenitors-specific chromatin remodeling complex(npBAF complex) and the neuron-specific	

chromatin remodeling complex(nBAF complex). During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to post-mitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes(nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth(By similarity). Has a strong influence on vitamin D-mediated transcriptional activity from an enhancer vitamin D receptor element(VDRE). May be a link between mammalian SWI-SNF-like chromatin remodeling complexes and the vitamin D receptor(VDR) heterodimer. Mediates critical interactions between nuclear receptors and the BRG1/SMARCA4 chromatin-remodeling complex for transactivation.

Molecular Weight:	58 kDa
Gene ID:	6602
UniProt:	Q96GM5

WB: 1:500-1:2000, IP: 1:200-1:1000, IHC: 1:20-1:200, IF: 1:20-1:200

Application Details

Application Notes:

• •	
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	PBS with 0.02 % sodium azide and 50 % glycerol pH 7.3,
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	-20°C for 12 months (Avoid repeated freeze / thaw cycles.)

	lI	1:
-	เวทก	111111
	land	11110

Expiry Date:

12 months