antibodies -online.com





PGK2 Protein (AA 1-417) (His tag)





Go to Product page

| _ | | | | | | |
|-----|---|---|----|----|-------------|---|
| () | V | ρ | rv | 16 | $\supset I$ | Λ |
| | | | | | | |

| Quantity: | 50 μg |
|-------------------------------|---|
| Target: | PGK2 |
| Protein Characteristics: | AA 1-417 |
| Origin: | Human |
| Source: | Escherichia coli (E. coli) |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This PGK2 protein is labelled with His tag. |
| Application: | Western Blotting (WB), ELISA |

Product Details

| Purity: | Greater than 95 % as determined by SDS-PAGE | |
|------------|---|--|
| Sterility: | 0.2 µm filtered | |

Target Details

| Target: | PGK2 | |
|-------------------|---|--|
| Alternative Name: | PGK2 (PGK2 Products) | |
| Background: | DJ417L20.2, PGK 2, pgk2, PGK2_HUMAN, PGKB, PGKPS, Phosphoglycerate kinase, Phosphoglycerate kinase 2, Phosphoglycerate kinase autosomal pseudogene, | |
| | Phosphoglycerate kinase testis specific, testis specific | |
| Molecular Weight: | 45.8 kDa | |

Target Details

UniProt: P07205


Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. |
|--------------------|--|
| Restrictions: | For Research Use only |

Handling

| Format: | Lyophilized |
|------------------|---|
| Reconstitution: | Centrifuge the vial at 10,000 rpm for 1 minute, reconstitute at 200 μ g/mL in sterile distilled water by gentle pipetting 2-3 times, don't vortex |
| Buffer: | Lyophilized from a 0.2 μ m filtered solution in 10 mM Hepes, 500 mM NaCl with 5 % trehalose, pH 7.4 |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | -20°C for 12 months as lyophilized, 2-8°C for 1 month under sterile conditions after reconstitution |
| Expiry Date: | 12 months |

Images



Western Blotting

Image 1.