antibodies.com

Datasheet for ABIN7125966 GPC1 Protein (Ser486Ala-Mutant, Ser488Ala-Mutant, Ser490Ala-Mutant) (His tag)



Overview

Quantity:	100 µg
Target:	GPC1
Protein Characteristics:	Ser486Ala-Mutant, Ser488Ala-Mutant, Ser490Ala-Mutant
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This GPC1 protein is labelled with His tag.

Product Details

Purpose:	Human Glypican 1 / GPC1 (S486A, S488A, S490A) Protein, His Tag (MALS verified)
Sequence:	Asp 24 - Ser 530
Characteristics:	Human Glypican 1 (S486A, S488A, S490A), His Tag is expressed from human 293 cells (HEK293). It contains AA Asp 24 - Ser 530 (Accession # P35052-1 (S486A, S488A, S490A).
Purity:	>95 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per μ g by the LAL method.
Grade:	MALS verified
Target Details	

Target:	GPC1
Alternative Name:	Glypican 1 / GPC1 (GPC1 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7125966 | 09/13/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
Background:	Synonyms: Glypican 1,GPC1,
	Glypican 1 has been shown to interact with SLIT2. This protein is involved in the misfolding of
	normal prion proteins in the cell membrane to the infectious prion form. Cell surface heparan
	sulfate proteoglycans are composed of a membrane-associated protein core substituted with a
	variable number of heparan sulfate chains. Members of the glypican-related integral membrane
	proteoglycan family (GRIPS) contain a core protein anchored to the cytoplasmic membrane via
	a glycosyl phosphatidylinositol linkage. These proteins may play a role in the control of cell
	division and growth regulation.
Molecular Weight:	57.9 kDa
NCBI Accession:	NP_002072
Pathways:	Glycosaminoglycan Metabolic Process, Regulation of Muscle Cell Differentiation
Application Details	
Application Notes:	This protein carries a polyhistidine tag at the C-terminus. The protein has a calculated MW of
	57.9 kDa. The protein migrates as 58-61 kDa under reducing (R) condition due to glycosylation.
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Buffer:	50 mM Tris, 150 mM NaCl, pH 7.5
Storage:	-20 °C
Storage Comment:	-20°C