Datasheet for ABIN7126086 Histone H2B Protein (full length) (rho-1D4 tag)

antibodies.com



Overview

Quantity:	0.5 mg
Target:	Histone H2B
Protein Characteristics:	full length
Origin:	CHO cells
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Histone H2B protein is labelled with rho-1D4 tag.
Application:	ELISA, Western Blotting (WB), SDS-PAGE (SDS), Crystallization (Crys), Functional Studies (Func)
Product Details	
Sequence:	MPEPSKSAPA PKKGSKKAIT KAQKKDGKKR KRSRKESYSV YVYKVLKQVH PDTGISSKAM
	GIMNSFVNDI FERIAGEASR LAHYNKRSTI TSREIQTAVR LLLPGELAKH AVSEGTKAVT KYTSSK
	Sequence without tag. The location of the tag depends on protein. You may also submit your
	preference when ordering.
Characteristics:	Made in Germany - from design to production - by highly experienced protein experts.
	CHO Histone H2B Protein (raised in Insect Cells) purified by multi-step, protein-specific
	process to ensure crystallization grade.State-of-the-art algorithm used for plasmid design (Gene synthesis).
	This protein is a custom-made protein and will be made for the first time for your order. This
	protein will be produced on the basis of on a Custom Service Project. We will make sure that
	every step in the production is successful from the design of the expression plasmid to the
	every step in the production is buobcostal norm the design of the expression plasmid to the

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7126086 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

	receive a correctly folded protein.
	The concentration of our recombinant proteins is measured using the absorbance at 280nm.
	The protein's absorbance will be measured in several dilutions and is measured against its
	specific reference buffer. The concentration of the protein is calculated using its specific
	absorption coefficient. We use the Expasy's protparam tool to determine the absorption
	coefficient of each protein.
Purification:	Three step purification of proteins expressed in baculovirus infected SF9 insect cells:
	1. Membrane proteins are fractioned by ultracentrifugation and subsequently solubilized with
	different detergents (detergent screen). Samples are analyzed by Western blot.
	2. The best performing detergent is used for solubilization and the proteins are purified via their
	rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate fractions are analyzed by Western blot.
	3. Protein containing fractions of the best purification are subjected to second purification step
	through size exclusion chromatograph. Eluate fractions are analyzed by SDS-PAGE and
	Western blot.
Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
Sterility:	0.22 µm filtered
Endotoxin Level:	Endotoxins have not been removed. Please contact us if you require an endotoxin-free version
	of this product.
Grade:	Crystallography grade
Biological Activity Comment:	Protein has not been tested for activity yet.

Target Details

Target:	Histone H2B
Abstract:	Histone H2B Products
UniProt:	G3HDU3

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	150 mM NaCL, 20 mM NaH2PO4 pH 7.4, 10 % glycerol. Note: Isoelectric point of protein taken into account regarding pH .
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)