

Datasheet for ABIN7126126

Recombinant anti-CDX2 antibody[Go to Product page](#)**4** Images

Overview

| | |
|----------------|--|
| Quantity: | 20 µg |
| Target: | CDX2 |
| Reactivity: | Human, Synthetic |
| Host: | Rabbit |
| Antibody Type: | Recombinant Antibody |
| Clonality: | Monoclonal |
| Conjugate: | This CDX2 antibody is un-conjugated |
| Application: | Immunohistochemistry (Formalin-fixed Sections) (IHC (f)) |

Product Details

| | |
|-----------------------------|---|
| Immunogen: | Recombinant fragments and synthetic peptides from human CDX2 protein (exact sequences are proprietary) |
| Isotype: | IgG |
| Specificity: | The intestine-specific transcription factors CDX1 and CDX2 are important for directing intestinal development, differentiation, proliferation and maintenance of the intestinal phenotype. CDX2 protein expression has been seen in GI carcinomas. Anti-CDX2 has been useful to establish GI origin of metastatic adenocarcinomas and carcinoids and is especially useful to distinguish metastatic colorectal adenocarcinoma from lung adenocarcinoma. However, mucinous carcinomas of the ovary also express CDX2 protein. It limits the usefulness of this marker in the distinction of metastatic colorectal adenocarcinoma from mucinous carcinoma of the ovary. |
| Cross-Reactivity (Details): | Human. |

Product Details

Purification: 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G.

Target Details


| | |
|-------------------|--|
| Target: | CDX2 |
| Alternative Name: | CDX2 (CDX2 Products) |
| Background: | Caudal type homeobox 2, Caudal type homeobox transcription factor 2, Caudal-type homeobox protein 2, CDX2,CDX2 / Caudal Type Homeobox 2 (GI Epithelial Marker) Cellular localisation: Nuclear |
| Molecular Weight: | 40kDa |
| Gene ID: | 1045, 174249 |
| UniProt: | Q99626 |
| Pathways: | Peptide Hormone Metabolism , Stem Cell Maintenance |

Application Details

| | |
|--------------------|--|
| Application Notes: | Positive Control: HT29 cells. Human colon carcinoma Known Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined. |
| Restrictions: | For Research Use only |


Handling

| | |
|--------------------|--|
| Concentration: | 200 µg/mL |
| Buffer: | Prepared in 10 mM PBS with 0.05 % BSA and 0.05 % azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-80 °C |
| Storage Comment: | Antibody with azide - store at 2 to 8 °C. Antibody is stable for 24 months. Non-hazardous. Also available WITHOUT BSA & azide at 1.0mg/ml. |




Western Blotting

Image 1. SDS-PAGE Analysis Purified CDX2 Recombinant Rabbit Monoclonal Antibody (CDX2/4394R). Confirmation of Purity and Integrity of Antibody.



Immunohistochemistry

Image 2. IHC analysis of formalin-fixed, paraffin-embedded human colon adenocarcinoma. Strong nuclear staining using CDX2/4394R at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2°: HRP-polymer, 30min. DAB, 5min.



Immunohistochemistry

Image 3. IHC analysis of formalin-fixed, paraffin-embedded human tonsil. Negative tissue control using CDX2/4394R at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2°: HRP-polymer, 30min. DAB, 5min.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN7126126.