

Datasheet for ABIN7126216

anti-CHEK2 antibody



Go to Product page

_					
	W	0	rv	10	W

Quantity:	100 μg	
Target:	CHEK2	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This CHEK2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunoprecipitation (IP), Flow Cytometry (FACS)	

Product Details

Immunogen:

Isotype: IgG2b Recognizes a serine/threonine protein kinase that is a required check-point to mediate cell cycle arrest, activation of DNA repair and apoptosis. In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. Also, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this	3 - 3 -	
cycle arrest, activation of DNA repair and apoptosis. In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. Also, this protein interacts with and	Isotype:	lgG2b
replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. Also, this protein interacts with and	Specificity:	Recognizes a serine/threonine protein kinase that is a required check-point to mediate cell
regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. Also, this protein interacts with and		cycle arrest, activation of DNA repair and apoptosis. In response to DNA damage and
tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. Also, this protein interacts with and		replication blocks, cell cycle progression is halted through the control of critical cell cycle
activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. Also, this protein interacts with and		regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative
blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. Also, this protein interacts with and		tumor suppressor. It contains a forkhead-associated protein interaction domain essential for
phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. Also, this protein interacts with and		activation in response to DNA damage and is rapidly phosphorylated in response to replication
suppressor protein p53, leading to cell cycle arrest in G1. Also, this protein interacts with and		blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C
		phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor
phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this		suppressor protein p53, leading to cell cycle arrest in G1. Also, this protein interacts with and
		phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this

Recombinant fragment of human CHEK2 corresponding to protein domain

Product Details

- Toddet Details			
	gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutations in TP53. Mutations in this gene are thought to confer a predisposition to sarcomas, breast cancer, and brain tumors.		
Cross-Reactivity (Details):	Human. Predicted to react with Mouse and Rat.		
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.		
Target Details			
Target:	CHEK2		
Alternative Name:	CHEK2 (CHEK2 Products)		
Background:	Cds1 homolog, Checkpoint kinase 2, Checkpoint like protein CHK2, CHK2 checkpoint homolog (S. pombe), hCds1, HuCds 1, LFS2, PP1425, RAD53, Rad53 homolog, Serine/threonine protein kinase Chk2,CHEK2 Cellular localisation: Nucleus.		
Molecular Weight:	60.91kDa		
Gene ID:	11200, 291363, 505297		
UniProt:	096017		
Pathways:	p53 Signaling, Apoptosis, Cell Division Cycle		
Application Details			
Application Notes:	Known_Application: Immunoprecipitation (1-2 μg per 100-500 μg of total protein), Flow Cytometry (1-2 μg/million cells), ,Western Blot (1-2 μg/mL),Immunofluorescence (1-2 μg/mL), ,Optimal dilution for a specific application should be determined. Positive_Control: HeLa or HCT 116 cells. Human testis, spleen or colon.		
Restrictions:	For Research Use only		
Handling			
Concentration:	1.0 mg/mL		
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.		
Preservative:	Azide free		
Storage:	-20 °C,-80 °C		

Handling

Storage Comment:	Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-	
	hazardous.	
Expiry Date:	24 months	