

Datasheet for ABIN7126312

anti-EIF2C3 antibody



Go to Product page

()	ve	r\/i		۱۸/
\cup	V C	1 / 1	$\overline{}$	٧V

Quantity:	100 μg
Target:	EIF2C3
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This EIF2C3 antibody is un-conjugated
Application:	ELISA, Immunoprecipitation (IP), Immunofluorescence (IF), Coating (Coat), Flow Cytometry (FACS)

Product Details

Immunogen:	Recombinant full-length human AGO3 protein
Isotype:	lgG1
Specificity:	This antigen is involved in binding and translation repression of short RNAs. This antigen is also
	known as argonaute RISC catalytic component 3. Required for RNA-mediated gene silencing
	(RNAi). Binds to short RNAs such as microRNAs (miRNAs) and represses the translation of
	mRNAs which are complementary to them. Proposed to be involved in stabilization of small
	RNA derivates (siRNA) derived from processed RNA polymerase III-transcribed Alu repeats
	containing a DR2 retinoic acid response element (RARE) in stem cells and in the subsequent
	siRNA-dependent degradation of a subset of RNA polymerase II-transcribed coding mRNAs by
	recruiting a mRNA decapping complex involving EDC4. Possesses RNA slicer activity but only
	on select RNAs bearing 5'- and 3'-flanking sequences to the region of guide-target
	complementarity.

Product Details Cross-Reactivity (Details): Human. Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G. **Target Details** Target: FIF2C3 Alternative Name: AGO3 (EIF2C3 Products) Background: Argonaute 3, Argonaute 3, elF 2C 3, elF-2C 3, elF2C 3, ElF2C3 protein, Eukaryotic translation initiation factor 2C 3, Eukaryotic translation initiation factor 2C3, FLJ12765, hAgo3, MGC86946, Protein argonaute-3, AGO3 (Argonaute-3) / elF2C3 Cellular localisation: Cytoplasm>P-bodies. Nucleoplasm. 97.36kDa Molecular Weight: Gene ID: 192669, 657659 UniProt: Q9H9G7 Fc-epsilon Receptor Signaling Pathway, Regulatory RNA Pathways, EGFR Signaling Pathway, Pathways: Neurotrophin Signaling Pathway **Application Details** Application Notes: Known_Application: ELISA (For coating, order antibody without BSA), ,Immunoprecipitation (1-2 μg per 100-500 μg of total protein (1 mL of cell lysate)), ,Flow Cytometry (1-2 μg/million cells), Immunofluorescence (1-2 µg/mL), ,Optimal dilution for a specific application should be determined. Positive_Control: HeLa, MCF7 or U-20S cells. Restrictions: For Research Use only

Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-

Prepared in 10 mM PBS, WITHOUT BSA and Azide.

Handling

Buffer:

Storage:

Concentration:

Preservative:

Storage Comment:

1.0 mg/mL

Azide free

-20 °C,-80 °C

Handling

	hazardous.
Expiry Date:	24 months