

Datasheet for ABIN7126420

anti-Fibronectin 1 antibody (AA 467-595)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	Fibronectin 1 (FN1)
Binding Specificity:	AA 467-595
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Fibronectin 1 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (Formalin-fixed Sections) (IHC (f)), Coating (Coat)

Product Details

Immunogen:	Recombinant fragment (around aa 467-595) of human fibronectin protein (exact sequence is proprietary)
Isotype:	IgG2b
Specificity:	Fibronectins are disulfide-linked, dimeric glycoproteins of ~440 kDa. They possess at least four binding sites for collagen, glycosaminoglycans, transglutaminase, and a cell surface receptor. Epitope of this MAb is located in the 2nd-3rd type-III repeats of fibronectin. Fibronectins are extracellular matrix glycoproteins that are essential for embryonic development. Fibronectins are also involved in cell adhesion, tissue organization, and wound healing. Fibronectins are present in basement membranes, interstitial connective tissue matrix, and blood. Cellular fibronectin is widely distributed in the stroma of many malignant tumors. This MAb reacts with human cellular fibronectin, but not plasma fibronectin.

Product Details

Cross-Reactivity (Details): Human.

Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target: Fibronectin 1 (FN1)

Alternative Name: FN1 ([FN1 Products](#))

Background: Cold insoluble globulin (CIG), FINC, FN1, FNZ, GFND, GFND2, LETS, Migration stimulating factor (MSF), Ugl-Y3,Fibronectin

Cellular localisation: Connective tissue matrix

Molecular Weight: 220kDa (monomer), 440kDa (dimer)

Gene ID: 2335, 203717

UniProt: [P02751](#)

Pathways: [Cellular Response to Molecule of Bacterial Origin](#), [Carbohydrate Homeostasis](#), [Autophagy](#)

Application Details

Application Notes: Known_Application: ELISA (For coating, order Ab without BSA),Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.

Positive_Control: SW156 cells. Human kidney.

Restrictions: For Research Use only

Handling

Concentration: 1.0 mg/mL

Buffer: Prepared in 10 mM PBS, WITHOUT BSA and Azide.

Preservative: Azide free

Storage: -20 °C,-80 °C

Storage Comment: Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.

Handling

Expiry Date: 24 months