

Datasheet for ABIN7126427 anti-Fibronectin 1 antibody (AA 467-595)



Overview

	100 μg	
Target:	Fibronectin 1 (FN1)	
Binding Specificity:	AA 467-595	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This Fibronectin 1 antibody is un-conjugated	
Application:	ELISA, Immunohistochemistry (Formalin-fixed Sections) (IHC (f)), Coating (Coat)	
Product Details		
Immunogen:	Recombinant fragment (around aa 467-595) of human fibronectin protein (exact sequence is	
	proprietary)	
Isotype:	IgG2a	
Specificity:	Fibronectins are disulfide-linked, dimeric glycoproteins of ~440 kDa. They possess at least four	
	binding sites for collagen, glycosaminoglycans, transglutaminase, and a cell surface receptor.	
	Epitope of this MAb is located in the 2nd-3rd type-III repeats of fibronectin. Fibronectins are	
	extracellular matrix glycoproteins that are essential for embryonic development. Fibronectins	
	extracellular matrix glycoproteins that are essential for embryonic development. Fibronectins are also involved in cell adhesion, tissue organization, and wound healing. Fibronectins are	
	are also involved in cell adhesion, tissue organization, and wound healing. Fibronectins are	
	Epitope of this MAD is located in the Zhu-sru type-III repeats of hibroriectin. Fibroriectins are	

Product Details Cross-Reactivity (Details): Human. Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G. **Target Details** Fibronectin 1 (FN1) Target: Alternative Name: FN1 (FN1 Products) Background: Cold insoluble globulin (CIG), FINC, FN1, FNZ, GFND, GFND2, LETS, Migration stimulating factor (MSF), Ugl-Y3, Fibronectin Cellular localisation: Connective tissue matrix Molecular Weight: 220kDa (monomer), 440kDa (dimer) 2335, 203717 Gene ID: UniProt: P02751 Cellular Response to Molecule of Bacterial Origin, Carbohydrate Homeostasis, Autophagy Pathways: **Application Details Application Notes:** Known_Application: ELISA (For coating, order Ab without BSA),Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined. Positive_Control: SW156 cells. Humankidney tissue. Restrictions: For Research Use only Handling Concentration: 1.0 mg/mL

Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-

Prepared in 10 mM PBS, WITHOUT BSA and Azide.

Azide free

-20 °C,-80 °C

hazardous.

Buffer:

Storage:

Preservative:

Storage Comment:

1 1		
\blacksquare	α	lina
	land	1111(
	I G I G	11119

Expiry Date:

24 months