

Datasheet for ABIN7126477

Recombinant anti-FOXP1 antibody (AA 350-450)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	FOXP1
Binding Specificity:	AA 350-450
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This FOXP1 antibody is un-conjugated
Application:	Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Immunogen:	Recombinant fragment (around aa350-450) of human FOXP1 protein (exact sequence is proprietary)
Isotype:	IgG
Specificity:	Transcriptional repressor. It plays an important role in the specification and differentiation of lung epithelium. Can act with CTBP1 to synergistically repress transcription but CTPBP1 is not essential. Essential transcriptional regulator of B cell development. FOXP1 protein plays an important role in the specification and differentiation of lung epithelium, also acts cooperatively with FOXP4 to regulate lung secretory epithelial cell fate and regeneration by restricting the goblet cell lineage program. FOXP1 is useful in subclassification of DLBCL and a high cutoff (80 %) for FOXP1 is needed to achieve high specificity for the ABC subtype.

Product Details

Cross-Reactivity (Details):	Human.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target:	FOXP1
Alternative Name:	FOXP1 (FOXP1 Products)
Background:	Fork head related protein like B, Forkhead box protein P1 (FOXP1), Glutamine rich factor 1, hFKH1B, HSPC215, QRF1,FOXP1 (Transcription Factor) Cellular localisation: Nucleus.
Molecular Weight:	75kDa
Gene ID:	27086
UniProt:	Q9H334
Pathways:	Chromatin Binding , Regulation of Muscle Cell Differentiation , Positive Regulation of Immune Effector Process , Production of Molecular Mediator of Immune Response

Application Details

Application Notes:	Known_Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined. Positive_Control: Human tonsil or colon.
Restrictions:	For Research Use only

Handling

Concentration:	1.0 mg/mL
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.

Handling

Expiry Date: 24 months