

Datasheet for ABIN7126923
anti-CCL8 antibody (AA 1-99)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	CCL8
Binding Specificity:	AA 1-99
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CCL8 antibody is un-conjugated
Application:	Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Immunogen:	Human recombinant MCP2 / CCL8 protein fragment (around aa1-99) (exact sequence is proprietary)
Isotype:	IgG1
Specificity:	This antimicrobial gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of N-terminal cysteine residues of the mature peptide. This chemokine is a member of the CC subfamily which is characterized by two adjacent cysteine residues. This cytokine displays chemotactic activity for monocytes, lymphocytes, basophils and eosinophils. By recruiting leukocytes to sites of inflammation this cytokine may contribute to tumor-associated leukocyte infiltration and to the antiviral state against HIV infection.

Product Details

Cross-Reactivity (Details):	Human.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target:	CCL8
Alternative Name:	CCL8 (CCL8 Products)
Background:	HC14, MCP2, MCP-2, SCYA8, SCYA10, Monocyte Chemotactic Protein 2 (MCP2) / CCL8 Cellular localisation: Secreted.
Molecular Weight:	11kDa
Gene ID:	6355, 271387
UniProt:	P80075

Application Details

Application Notes:	Known_Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined. Positive_Control: Human small intestine, spleen or adrenal gland tissue.
Restrictions:	For Research Use only

Handling

Concentration:	1.0 mg/mL
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.
Expiry Date:	24 months