

Datasheet for ABIN7127074

Recombinant anti-Calretinin antibody (AA 23-242)



Overview

Quantity:	100 μg
Target:	Calretinin (CALB2)
Binding Specificity:	AA 23-242
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This Calretinin antibody is un-conjugated
Application:	Immunohistochemistry (Formalin-fixed Sections) (IHC (f))
Product Details	
lmmunogen:	Recombinant human Calretinin (Calbindin 2) protein fragment (around aa23-242) (exact sequence is proprietary)
Isotype:	IgG
Specificity:	It recognizes a protein of about 29 kDa, which is identified as Calretinin (also known as Calbindin 2). Calretinin is a vitamin D-dependent calcium-binding protein involved in calcium signaling. It is present in subsets of neurons throughout the brain and spinal chord, including sensory ganglia. Antibody to calretinin is useful in differentiating mesothelioma from adenocarcinomas of the lung. It also aids in differentiating adrenal cortical neoplasms from pheochromocytomas.
Cross-Reactivity (Details):	Human.

Product Details Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G. Target Details Target: Calretinin (CALB2) Alternative Name CALB2 (CALB2 Products) Background: 29 kDa calbindin, CAB29, CAL2, CALB2, Calbindin 2, Calbindin D29K, Calretinin, CR, Calretinin / Calbindin 2 (Mesothelioma Marker) Cellular localisation: Cytoplasm. Nucleus. Molecular Weight: 29kDa Gene ID: 794, 106857 UniProt: P22676 **Application Details** Application Notes: Known_Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined. Positive_Control: Human brain, testis or mesothelioma tissue. Restrictions: For Research Use only Handling Concentration: 1.0 mg/mL Buffer: Prepared in 10 mM PBS, WITHOUT BSA and Azide. Preservative: Azide free Storage: -20 °C,-80 °C Storage Comment: Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-

hazardous.

24 months

Expiry Date: